

Ten lessons for controlling invasive species: Wisdom from the long-standing sea lamprey control program on the Laurentian Great Lakes

Steven J. Cooke , Carrie L. Baker, Julia L. Mida Hinderer, Michael Siefkes, Jessica M. Barber, Todd B. Steeves, Margaret F. Docker, Weiming Li, Michael P. Wilkie, Michael L. Jones, Kelly F. Robinson, Erin S. Dunlop, Cory O. Brant, Nicholas S. Johnson, William P. Mattes, Marc Gaden and Andrew M. Muir

Steven J. Cooke (steven_cooke@carleton.ca) is affiliated with the Department of Biology and Institute of Environmental and Interdisciplinary Science at Carleton University, in Ottawa, Ontario, Canada. Carrie L. Baker, Julia L. Mida Hinderer, Michael Siefkes, Marc Gaden, and Andrew M. Muir are affiliated with the Great Lakes Fishery Commission, in Ann Arbor, Michigan, in the United States. Jessica M. Barber is affiliated with the US Fish and Wildlife Service, Marquette Biological Station, in Marquette, Michigan, in the United States. Todd B. Steeves is affiliated with the Canadian Department of Fisheries and Oceans, in Sault Ste Marie, Ontario, Canada. Margaret F. Docker is affiliated with the Department of Biological Sciences at the University of Manitoba, in Winnipeg, Manitoba, Canada. Weiming Li and Michael L. Jones are affiliated with the Department of Fisheries and Wildlife at Michigan State University, in East Lansing, Michigan, in the United States. Michael P. Wilkie is affiliated with the Department of Biology and Laurier Institute of Water Science at Wilfrid Laurier University, in Waterloo, Ontario, Canada. Kelly F. Robinson is affiliated with the US Geological Survey, Georgia Cooperative Fish and Wildlife Research Unit and with the Warnell School of Forestry and Natural Resources at the University of Georgia, in Athens, Georgia, in the United States. Erin S. Dunlop is affiliated with the Aquatic Research Section at the Ontario Ministry of Natural Resources, at Trent University, in Peterborough, Ontario, Canada. Cory O. Brant is affiliated with the US Geological Survey Great Lakes Science Center, in Ann Arbor, Michigan, in the United States. Nicholas S. Johnson is affiliated with the US Geological Survey, Great Lakes Science Center, Hammond Bay Biological Station, in Millersburg, Michigan, in the United States. William P. Mattes is affiliated with the Great Lakes Indian Fish and Wildlife Commission, in Odanah, Wisconsin, in the United States.

Abstract

Sea lamprey (*Petromyzon marinus*) control in the Laurentian Great Lakes of North America is among the largest and most successful control programs of an invasive species anywhere on the planet. The effort began more than 75 years ago; it unites multiple nations, states, and provinces with the common goal of controlling this invasive species and protecting a valuable fishery. The science-based control program is administered by the Great Lakes Fishery Commission (GLFC), a body arising from a treaty signed by the United States and Canada. In the present article, we share 10 lessons learned from decades of successful sea lamprey control with the hopes of informing ongoing and future control programs targeting biological invasions. The 10 lessons we identified are to act boldly in times of crisis, to maintain the social license, to invest in capacity building, to break down the silos, to support fundamental science, to diversify your portfolio of control measures, to strive for continuous improvement, to confront the trade-off between information and action, to keep your foot on the gas, and to keep your eyes on the prize. The GLFC has long fostered a framework that uses some military strategy and verbiage that extends across the lessons (e.g., know your enemy). Other lessons are more nascent as the GLFC reenvision its relationship with Indigenous peoples and governments in a path to reconciliation where two-eyed seeing is being embraced. Through adaptive management, horizon scanning methods, and embracing implementation science, the lessons learned about sea lamprey control will continue to evolve, which is itself a lesson. We submit that the lessons shared in the present article will help guide invasive species control programs spanning taxa, ecosystems, and regions.

Keywords: invasive species, environmental science, conservation

Invasive species represent a major threat to biodiversity (Mack et al. 2000, McGeoch et al. 2010, IPBES 2019) and are considered a key driver of population declines (e.g., Bellard et al. 2016) and even extinctions (Bellard et al. 2021). The impacts of invasions extend beyond individual populations and species, often resulting in dramatic changes in ecosystem structure and function (Crystal-Ornelas and Lockwood 2020). The economic costs of invasive species are immense (Diagne et al. 2021, Ahmed et al. 2023) and eclipse any benefits that invasions may yield (Cameiro et al. 2024). In fact, the effects of biological invasions have manifold impacts on ecosystem services (Vilà and Hulme 2017) extending to human wellbeing and food security (Linders et al. 2020, Rai and Singh 2020). For the aforementioned reasons, there are many efforts to prevent biological invasions (Mack et al. 2000, Keller et al. 2008) ranging from biosecurity initiatives that restrict trade

and transfer of invasive species using regulatory mechanisms (for a review, see Lieurance et al. 2023) to educational efforts intended to engage the public in such efforts (for a review, see Haley et al. 2023). Such prevention methods are the obvious first line of defense (Briski et al. 2013) and are often substantially cheaper than postinvasion control (Leung et al. 2002). The management of biological invasions sometimes relies on a rapid response approach, with a focus on eliminating organisms before establishment and, in some cases, the need for comprehensive control programs (Burgeil 2020, Reaser et al. 2020, Herbst et al. 2021).

Despite a focus on prevention, biological invasions do occur, which, in some instances, necessitates invasive species control measures to be used or control programs to be developed (Mack et al. 2000). However, once invasive species become established, control may be challenging or, more often, impossible. For

Received: March 21, 2025. Revised: July 16, 2025. Accepted: July 28, 2025

© The Author(s) 2025. Published by Oxford University Press on behalf of the American Institute of Biological Sciences. This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact reprints@oup.com for reprints and translation rights for reprints. All other permissions can be obtained through our RightsLink service via the Permissions link on the article page on our site—for further information please contact journals.permissions@oup.com




Figure 1. Schematic of the life cycle of sea lamprey. Spawning stage: (1a) Sea lampreys use their sucker mouth to suction onto rocks and make crescent-shaped nests for spawning. (1b) Eggs are deposited in the nests by females while being fertilized by males. Larval stage: (2a) About two weeks after fertilization, larvae hatch from the eggs. Lacking eyes and a sucker mouth, harmless larvae drift downstream and settle into a silt or sand stream bottom. (2b) Larvae live in stream bottoms for 3–10 years depending on conditions. Larvae feed by filtering organic particles, algae, and microscopic organisms from the sediment and water. Metamorphosis: (3) Once large enough, larvae metamorphose, growing sucker mouths, teeth, and eyes. (4) At the completion of metamorphosis, sea lampreys emerge from the stream bottom. Postmetamorphosis: (5) The newly metamorphosed parasitic juveniles migrate downstream into large rivers and lakes where they feed on fish (as parasites) until it is time to spawn. After feeding for 1–1.5 years, the sea lamprey's digestive system shuts down and they become sexually mature. (6) They migrate into streams to spawn and then die, completing the life cycle.

instances where biological invasions create extreme levels of negative impacts on ecosystem services, a coordinated control program may need to be sustained for long periods (Simberloff 2014). In this context, control is defined as efforts intended to reduce the abundance and impacts of a given invader. We intentionally use that term given the inherent challenges with fully eradicating invaders, which may not be needed to reduce impacts to reasonable levels (Bomford and O'Brien 1995). Examples of prominent invasive species control programs can be found around the globe and include the cane toad (*Bufo marinus*) in Australia (Shanmuganathan et al. 2010), the brown tree snake (*Boiga irregularis*) in Guam (Hall 1996), lake charr (trout) *Salvelinus namaycush* in Yellowstone Lake (in the United States; Koel et al. 2020), prickly pear cacti (*Opuntia* spp.) particularly in Australia and South Africa (Humphries et al. 2022), and Asian longhorn beetles (*Anoplophora glabripennis*) in Europe and North America (Haack et al. 2010). Invasive species control programs can be operational for extended periods (i.e., decades) and are often supported by extensive research and monitoring. Although a vast body of research examines the effectiveness of invasive species control initiatives, much of it is focused on the biological aspects (e.g., which removal methods work and in which contexts; for a review, see Metha et al. 2007 and Coleman et al. 2017) but does not consider the institutional and procedural aspects of control programs that underpin the science-based control efforts.

One of the longest-standing control programs for an invasive species is that of sea lamprey (*Petromyzon marinus*; see figure 1 for life cycle) in the Laurentian Great Lakes of North America (see box 1). The Sea Lamprey Control Program is coordinated by the Great Lakes Fishery Commission (GLFC) as part of a treaty signed by Canada and the United States in 1954. Although eradication is currently improbable, control efforts have suppressed sea lamprey populations by about 90% (Siefkes et al. 2021), allowing native fish populations and introduced recreational fishes to thrive (Robinson et al. 2021a, Wingfield et al. 2021).

Given that the sea lamprey control program has deployed control efforts on the Laurentian Great Lakes for more than 65 years (and is therefore one of the longest-running and largest of such programs) and has explored control methods for over 75 years, the program provides a unique opportunity to reflect on successes and failures. In the present article, we present lessons learned over more than a half century of invasive sea lamprey control with the hopes of informing ongoing and future control programs targeting other biological invasions. Billions of dollars have been spent globally on invasive species control measures and programs (Jardine and Sanchirico 2018), reinforcing the need for such reflections. In the present article, we intentionally focus on conceptual aspects related to control program delivery that should be readily transferable to other contexts rather than dwelling on specific control measures. This article is novel in that, to our knowledge, there are few attempts to consider the broader context in which invasive species control programs succeed or fail. Several other synthetic accounts of the sea lamprey control story exist (see Brant 2019, Brant et al. 2019, McLaughlin et al. 2021, Wingfield et al. 2021), but none of these has been framed in terms of transferrable lessons for controlling other invaders. Although we attempt to anchor each of our lessons in empirical literature, the reality is that the majority of these lessons arise from frontline efforts and behind-the-scenes support that collectively enable the sea lamprey control program but have not been well described in the literature. The lessons in the present article are not intended to be prescriptive but rather serve as ideas, inspiration, and candid advice (based on successes and failures) for others working on the control of biological invasions. Similar reflective, lessons-oriented papers related to coordinated invasive species control are lacking (but see Dahlberg et al. 2023 for a more focused retrospective analysis of Dreissenid mussel control efforts in open waters and more general perspectives in Myers et al. 2000, Simberloff 2009, and Phillips et al. 2019 or regional overviews such as Davies et al. 2020). To our knowledge, the present article is unique in its

Box 1. An overview of lamprey control measures.

The present control program has a 65-year history based on sound science, dogged determination, collaborative partnerships, and a little luck. Little was known about sea lamprey when they invaded the Great Lakes, in particular their life history (figure 1). Research and experience revealed that sea lamprey are most effectively controlled in streams, either during their spawning migration as mature adults or during their in-stream residence as larvae. Initial efforts relied on crude electrical barriers to halt migration (figure 2a),




Figure 2. (a) Historical electric barrier used to guide lamprey. (b) TFM treatments occur in lotic systems (e.g., rivers, streams) targeting larval stage lamprey. Photograph: Zak Allen. (c) Granular bayluscicide treatments are conducted using boats to target larval lampreys in more lentic (e.g., lake) environments. Photograph: Zak Allen. (d) Low-head barriers are installed on tributaries to prevent upriver migrating lamprey from reaching spawning grounds. Photograph: Ted Lawrence. (e) Traps are used often alongside barriers to capture adult lamprey for assessment and euthanasia. Photograph: Marc Gaden. All of the photos part of the GLFC archive and are used with permission.

but the method was dangerous and nonselective, and ineffective at a control scale (Brant 2019). Through a series of rigorous and persistent tests, researchers found a chemical called 3-trifluoromethyl-4-nitrophenol (TFM), that was selective for lampreys, environmentally benign, and safe for human handling (figure 2b). Although TFM can be toxic to a few nontarget species, such as juvenile lake sturgeon *Acipenser fulvescens* (more than 100 millimeters) and mudpuppy *Necturus maculosus* under certain

(Continued)

Box 1. An overview of lamprey control measures.

environmental conditions, it has become the cornerstone of sea lamprey control. A second chemical, 2',5-dichloro-4'-nitrosalicylanilide, or Bayluscide, is used as an additive to enable more judicious use of TFM in large rivers and by itself in lentic areas such as embayments adjacent to river mouths (figure 2c). Barriers to migration, such as purpose-built low-head sea lamprey barriers and dams constructed for other purposes, remain an effective tool to limit sea lamprey distribution and access to spawning habitat (figure 2d; Zielinski et al. 2019). Sea lamprey have infested around 10% of Great Lakes tributaries, and extensive larval assessment operations serve to determine the abundance and distribution of sea lamprey larvae in those tributaries, thereby prioritizing use of limited control resources. Finally, a network of assessment traps deployed in tributaries to intercept migrating adults (figure 2e) provides opportunity to understand relative changes in sea lamprey abundance through time, remove migrating sea lamprey to prevent spawning, and to provide sea lamprey for outreach and ongoing research to advance sea lamprey control techniques. Those aforementioned methods are variously supplemented (see Siefkes et al. 2021) by portable electric barriers, juvenile trapping, sea lamprey sterilization (currently on operational pause), and pheromone or alarm cues (to attract or repel lamprey to improve the effectiveness of other control methods). Sea lamprey control costs roughly US\$25 million a year to protect one of the single largest freshwater ecosystems on earth, with a recreational fishery estimated to be worth US\$5.1 billion annually (Cornicelli et al., 2022).




Figure 3. (a) Oral disc of sea lamprey used for attachment to the fish (via keratinized teeth) prior to a rasping tongue being used to feed. (b) Fish with lamprey scar. (c) Lampreys engaging in spawning behavior. (d) Sea lamprey form. Photographs: Andrew Muir.

focus on a coordinated control program targeted toward a single invader.

Ten lessons

The 10 lessons described below were identified through an iterative process where the first step involved building a diverse team. The present coauthors collectively represent scientists (from different disciplines and realms including academia, Indigenous governments, and federal governments), practitioners (including sea lamprey control agents, which we call *control agents*—Fisheries and Oceans Canada and the US Fish and Wildlife Service staff that deliver sea lamprey control on behalf of the GLFC), and GLFC technical, science transfer, policy, and leadership staff. All of the present coauthors have participated in the control program in various ways including through service on various GLFC committees and boards (e.g., the Sea Lamprey Control Board, the Sea Lamprey Research Board). Collectively, the coauthors have more than 275 years of experience with lamprey control. Ideas for lessons were collated, thematized, and where necessary merged or split, to create a final list of 10 lessons. Given the diversity in training, experience, and perspective of our author team, we are confident that the lessons identified in the present article are indeed the

key lessons that have emerged from the Sea Lamprey Control Program and are potentially relevant to other control programs.

Act boldly in times of crisis

The control program arose out of crisis. By 1940, when sea lampreys had made it to all five Great Lakes, native piscivorous fishes, such as the lake charr, and their prey were already dramatically reduced in abundance (Muir et al. 2012). Stakeholders and rightsholders, such as commercial fishers, business owners, Indigenous or tribal community members, and recreationalists, witnessed the destruction firsthand when they saw fish with sea lamprey wounds (see figure 3), experienced sharply reduced fish harvest, and as a result of release from predation pressure by lake charr, dealt with beaches littered with dead invasive alewives (*Alosa pseudoharengus*) that had exploded in abundance (Smith 1968, Gaden et al. 2022). The crisis of collapsed fisheries and an unknown invader provided the perfect opportunity to rally jurisdictions and politicians to overcome the impediments to nine previously failed attempts to establish interjurisdictional cooperation in the basin. Failure of individual jurisdictions to control sea lampreys on their own was seized as the fuel for finally establishing cooperative fishery management in the Great Lakes (Gaden et al. 2022). Stakeholders in both Canada and the United States, noting the destruction of valued fisheries, persuaded politicians

Box 2. Two-eyed seeing and sea lamprey control.

Two-eyed seeing, or *etuaqtmumk* (a Mi'kmaw word and concept), is learning to see with the strengths of Indigenous ways of knowing together with the strengths of western science, in order to gain a broader perspective for the benefit of all (Bartlett et al. 2012). Although this approach had its beginnings in the health field, it is now being increasingly used in fisheries research, management, and stewardship contexts (Almack et al. 2023, Duncan et al. 2023, Reid et al. 2021). Although one US treaty organization, the Great Lakes Indian Fish and Wildlife Commission (1842 Treaty waters) is represented on the Sea Lamprey Control Board (SLCB; www.glfc.org/sea-lamprey-control-board-task-forces.php), the control program has not yet used two-eyed seeing to guide the decision-making process, a shortcoming considering the many Tribes and First Nations who have inherent rights to the lands and waters of the Great Lakes basin (Mattes and Kitson 2021, Nonkes et al. 2024). However, efforts are underway to better understand Indigenous perspectives of sea lamprey and the control program and to include this previously overlooked source of expertise and wisdom within the sea lamprey research and control programs (Nonkes et al. 2023). Adopting a two-eyed seeing approach is crucial not only in the pursuit of reconciliation but also considering the potential declines in social acceptance of sea lamprey control among Indigenous communities as well as the value added by looking at the issue from a different perspective.

to establish the GLFC by treaty to lead a multinational control effort and, importantly, to fund the operations (Gaden et al. 2022). Applied science to combat the sea lamprey was prioritized and, to this day, effective cross-jurisdiction resource sharing, coordinated by the GLFC, remains critical for the success of the control program (Brant et al. 2019, Gaden et al. 2022). We acknowledge that, during the early phases of an invasion, acting boldly may benefit from being done with expediency in an effort to eradicate an invader before getting to the point where ongoing control is necessary.

Maintain the social license

The control program has been a tremendous success, with sea lamprey populations reduced by at least 90% in most areas of the Great Lakes basin (Siefkes 2017). However, although a wide range of stakeholders and politicians continue to support the control program, the worst effects of invasive sea lamprey haven't been witnessed in 60 years (but see lesson no. 9), so the perceived risk of sea lamprey, and the continuing social acceptance for the control methods are not fully appreciated (Gaden et al. 2021b). The loss of the social license for sea lamprey control could range from limitations on the type of controls that can be used (including challenges with permitting) or, at worst, termination of the control program. The two main techniques for sea lamprey control, pesticides and dams, might not be as socially acceptable as they have been in the past (e.g., pesticides represent addition of chemicals to waters used by people and can result in effects on nontarget species, such as native Unionid mussels, impacts, dams can block native species and cause human safety concerns; Gaden et al. 2021b). Although the GLFC is exploring means of maintaining the social license for sea lamprey control, practitioners must maintain an active effort to provide policymakers and the public with timely, relevant, and factual information about sea lamprey, their effects, and their control. The GLFC and its partners regularly engage with a range of stakeholders, rightsholders, and politicians to keep people informed about the benefits and results of the control program. Likewise, legislative staff from the GLFC work on a regular basis with policymakers and elected officials to demonstrate the importance of invasive species control and to provide data about the economic benefits of sea lamprey control. This communications effort is a two-way street, allowing for the GLFC and its partners to convey the social, ecological, and economic benefits of the control program but also to hear and understand public concerns about the program. Outreach includes near-constant delivery of clear and consistent

messaging through social media, web pages, videos, and printed materials, as well as many in-person, interactive venues featuring live lamprey. Another ongoing concern is how changing government policy and funding shortfalls could undermine invasive species control or eradication programs, such as in Australia, where funding cuts could threaten the success of an invasive fire ant (*Solenopsis invicta*) eradication plan (Wylie and Janssen-May 2017). In recognition of the moral and legal imperative to engage Indigenous communities and governments, the GLFC has begun to build better relationships with Indigenous nations and embrace a two-eyed seeing approach (see Reid et al. 2021), which strives for learning to see with the strengths of both Indigenous and Western ways of knowing (box 2; Bartlett et al. 2012). This element of the GLFCs evolution is too novel to consider it a lesson (see box 2 for early learnings), but it underpins the importance of self-reflection and the ability to redress past inadequacies in the delivery of its programs.

Invest in capacity building

Investing in capacity building is critical for the success and longevity of a control program. Capacity building is a process of enhancing the skills of individuals, organizations, and communities, leading to improved program effectiveness, sustainability, and innovation (Eade 2007, Sobek and Agius 2007, Labin et al. 2012). It also promotes collaboration among management agencies, Indigenous stewards, and researchers, which is vital for mutual learning to inform action (Wiener et al. 2011, Jellinek et al. 2021). Capacity building partnerships capitalize on investments by leveraging multiple resources and enhancing communication leading to collaborative decision-making and better stewardship (Behnken et al. 2016). The GLFC has fostered the training and development of students, early career scientists, and control agents, many of whom have gone on to devote lengthy careers to sea lamprey control in either research or management. It is also important to diversify (Mullin et al. 2021), because diverse teams are better at problem solving (Cheruvilil et al. 2014), ultimately driving positive change and improving control program delivery. To this end, the GLFC is exploring ways to break down barriers to access members of groups that are currently underrepresented in the fisheries field, with an ultimate goal of fostering a more diverse and therefore effective workforce to carry out sea lamprey research and management into the future.

Break down the silos

The control program minimizes siloed thinking by efficiently communicating and coordinating across its own elements and with resource management agencies and science institutions throughout the Great Lakes. Owing to the coordination elements of its mandate, the GLFC serves as a knowledge broker, representing a fundamental central node in the broader Great Lakes environmental governance network (Mulvaney et al. 2015, Song et al. 2019, Song et al. 2020). The control program also adopts and integrates new science and technology through the coproduction and transfer of science to decision-makers (Gaden et al. 2021a and in the present article). Collaboration within the control program is facilitated through the SLCB, which advises the GLFC in implementing control strategies, policies, and programs. The SLCB membership includes representatives from federal, state, provincial, and tribal fishery agencies, along with nongovernmental and academic institutions, and importantly includes both researchers and managers. Coordination with resource management agencies throughout the Great Lakes occurs through GLFC-sponsored lake technical committees (www.glfc.org/joint-strategic-plan-committees.php), of which sea lamprey control agents are invited members on each Great Lake. The SLCB is advised by control program task forces, composed largely of technical folks who advise on the science priorities (Krueger and Marsden 2007). The SLCB research priorities drive the focus of the GLFCs research program. This feedback loop between science and management ensures adherence to the mission. Research proposals are evaluated against the priorities by the Sea Lamprey Research Board (SLRB; <https://www.glfc.org/sea-lamprey-research-board.php>), which consists of both scientific experts and control agents. Funded research is often coproduced with control program staff (Lewandoski et al. 2021, McLaughlin et al. 2021) and the GLFC sponsors the Science Transfer Board (<https://www.glfc.org/science-transfer-board.php>), which further facilitates knowledge exchange between scientists and control program staff (Hinderer et al. 2021). Embedding scientists within implementation-focused groups (e.g., the SLCB) and embedding sea lamprey control staff within science-focused programs (e.g., the SLRB) also helps bridge the knowledge-action gap (Nguyen et al. 2017) and promotes evidence-based decision-making resulting in more effective sea lamprey control by providing a direct pipeline for new science into control policies and operating procedures. Therefore, the GLFC has strategically worked to break down the silos through strong communication and establishing and nurturing fora for integration, knowledge sharing, coproduction, and consensus decision-making.

Support fundamental science

Although it is focused on research to augment or improve control methods, the Sea Lamprey Research Program (SLRP) has also played the long game by investing in fundamental science. Fundamental science serves as the foundation for more mission-directed research and may pay unexpected dividends. Sea lamprey pheromone research started in sensory neurobiology in the 1950s and 1960s (see Buchinger et al. 2015); subsequent investments in physiological and behavioral research (e.g., Li et al. 1995, Daghfous et al. 2016), and the identification of the active pheromone components (Li et al. 2018) and their receptors and antagonists (Zhang et al. 2020, Scott et al. 2023) has led to the potential to use pheromones to disrupt sea lamprey migration or spawning. Likewise, a 40-year quest to elucidate the hypothalamic-pituitary system in lampreys, while shedding

light on the evolution of this uniquely vertebrate feature, may eventually enable methods to suppress sea lamprey reproduction (Sower 2015, 2018). Similarly, sea lamprey genome sequencing and evolutionary developmental biology studies date back to a long-term interest in this jawless vertebrate as a model to recapitulate the early evolution of vertebrates (York et al. 2019) and the vertebrate genome (Smith et al. 2013, 2018). These genomic resources will enable research into possible genetic control of sea lamprey (Ferreira-Martins et al. 2021). A better understanding of the sea lamprey fundamental biology has yet to pay dividends in terms of broadscale control but has laid the groundwork for advancing novel control tools with a different mechanism of action than TFM (Wilkie et al. 2019), advancing selective fish passage whereby sea lampreys are selectively removed from a migrating stream of fishes (Zielinski et al. 2020) and elucidating the mechanism of sex determination and potential gene targets for genetic control (McCauley et al. 2015).

Diversify your portfolio of control measures

The control program adopted an integrated approach to controlling sea lamprey in the 1980s (Sawyer 1980) and has since invested roughly 50% of the annual research budget to developing new control tools (Siefkes et al. 2021). Tactics developed to target the adult stage to reduce reproduction include direct removal of adults in traps with trap encounter rates increased with physical leads and electrical, chemosensory, light, and auditory stimuli (Miehls et al. 2020) and the reduction in spawning success through the release of sterilized males (Johnson et al. 2024) or chemosensory cues that disrupt mate finding (Scott et al. 2023). Tactics have also been developed to remove juvenile sea lamprey during their outmigration from rivers to lakes (i.e., those not killed by lampricide treatment that metamorphose into the harmful parasitic stage). Juvenile sea lamprey encounter rates with traps can be improved using physical diversion, electrical repellents, chemosensory repellents, and lights (Evans et al. 2021). The control program recently reemphasized how new or supplemental control tools can diversify the suite of available tactics and potentially reduce reliance on lampricides (Siefkes et al. 2021), which is needed to prevent or slow the development of pesticide resistance (Dunlop et al. 2018). This renewed supplemental control philosophy is currently being implemented in a framework designed to enhance learning through adaptive management and tailor tactics to align with the environmental and social context (Lewandoski et al. 2021, 2025).

Strive for continuous improvement

Invasive species control requires making decisions in the face of uncertainty (Johnson et al. 2017, Robinson et al. 2021b). Therefore, an adaptive management approach (Walters 1986), in which a reduction in uncertainty leads to refinement of actions and, at times, a complete course change is necessary for success. This requires striving for continuous improvement. Indeed, failure and surprise are expected in adaptive management programs, requiring flexibility to seize them as learning opportunities (Allen and Gunderson 2011). Inspired by the first Sea Lamprey International Symposium (see *Canadian Journal of Fisheries and Aquatic Sciences* vol. 37, issue 11), over the last 45 years, the control program has maintained and enhanced its flexibility in response to learning, as well as a great deal of social and environmental change. This includes learning through experimentation and adaptive management principles but also through failure and surprise (e.g., Sullivan et al. 2021). Some of the first attempts to control sea lamprey did not have the desired or anticipated outcomes, but the

flexibility to explore supplemental control options has been effective and has been embraced as part of a new adaptive management program (Lewandoski et al. 2025). Indeed, learning from failure has served several rodent control programs well (Holmes et al. 2015, Samaniego et al. 2021). The coordination of the control program, multiple task forces, the SLRB, and many councils and committees allows for feedback among control actions, monitoring, and targeted research. In addition, the trust built through these long-term collaborations among agencies and between countries, facilitated by the GLFC, over the course of the implementation of sea lamprey control has made this programmatic flexibility possible (van Putten et al. 2022, Muir et al. 2023). Continual control program assessment and associated investments in implementation science (Cooke et al. 2024) allow evidence-informed refinement to occur such that there is opportunity for continuous improvement.

Confront the trade-off between information and action

Since the early stages of sea lamprey control, the control program has relied on lampricides to remove larval sea lampreys before they metamorphose and become parasites. The larval stage typically lasts 3–5 years, but the time to metamorphosis can vary among streams and over time (Brant 2019). In a control program where treatment resources are limited, treating streams with lampricides earlier than necessary (i.e., years before metamorphosis) presents financial and opportunity costs while treating too late allows escapement of large numbers of parasites. The Great Lakes basin is expansive (approximately 240,000 square kilometers) and contains hundreds of spawning streams where sea lampreys flourish, so the GLFC has used larval assessment to prioritize streams for annual treatments. Larval assessment is costly, and there is a trade-off between doing more assessment or allocating resources to treat more streams. Using a field-tested adaptive management approach (Hansen and Jones 2008a) and simulation models (Jones et al. 2009), researchers determined that a less costly and precise assessment approach actually led to better outcomes (i.e., greater suppression for the same cost; Jubar et al. 2021). In addition, control agents realized many streams reliably produced new populations at a predictable time after treatment, leading to the establishment of expert judgment streams requiring even less assessment and gaining further efficiencies. This illustrates the use of a value-of-information approach to guide decision-making, an approach with broad, and underappreciated, salience to fisheries management (Hansen and Jones 2008b).

Keep your foot on the gas

A knowledge of when larvae were likely to metamorphose, and rigorous sea lamprey assessment to prioritize which streams require treatment (Hansen and Jones 2008a), yielded unprecedented success in suppressing sea lamprey in the mid- to late 2000s (Sullivan et al. 2021). However, the need for managers to keep their foot on the gas was made clear during the COVID-19 pandemic of 2020–2022, which demonstrated that even a temporary disruption to the control program could undermine success (Marcy-Quay et al. 2025). The lockdowns and travel restrictions due to COVID-19 in 2020 greatly curtailed sea lamprey assessments and lampricide treatments, with only 26 of 101 scheduled streams undergoing lampricide treatments in 2020 (Burkett et al. 2021). Those reductions resulted in a spike in lamprey abundance and concomitant increase in fish wounding in Lakes Superior and Huron in 2021 compared with 2019 (Marcy-Quay et al. 2025). In 2022, when the bulk of metamorphosed sea lamprey from 2020

had likely matured, the returning adult sea lamprey populations were well above their abundance targets in all five Great Lakes (Marcy-Quay et al. 2025). The cost to the ecosystem was stark; estimates indicated that there were more than 50,000 additional parasitic sea lampreys produced in the five Great Lakes in 2022, amounting to upward of 1,000,000 kilograms of fish killed, based on prior estimates that a single sea lamprey can kill approximately 18 kilograms of fish during its parasitic phase (Swink 2003). It remains to be seen how quickly sea lamprey populations will be brought back under control in coming years, but there is reason for optimism as the control program remains well-funded and normal operations have resumed. The COVID-19 pandemic provides a cautionary case study that invasive species control measures cannot be relaxed, even temporarily, which is salient given changing priorities associated with transitions in leadership, dynamic geopolitical forces, uncertain financial markets, and associated variation in funding for control programs.

Keep your eyes on the prize

Sea lamprey control is a means to a more fundamental set of objectives in the Great Lakes basin, the most notable of which is maintaining sustainable lake charr populations for recreational and commercial fisheries (Gaden et al. 2021b). Over time, these objectives have diversified to include other ecological, economic, and social goals, such as healthy populations of desired species, recreational and commercial user satisfaction, and a lucrative economy. By defining and refining a set of achievable fundamental objectives, the control program remains focused on ultimate needs of the social-ecological system. In addition, these fundamental objectives reach beyond sea lamprey control, providing a basis for other decisions in fisheries management, habitat restoration, and native species restoration that are ongoing in the Great Lakes basin. As a knowledge brokering and boundary organization, the GLFC successfully coordinates sea lamprey control and fishery management across these multiple objectives, jurisdictions, and organizations. Looking to the future, double- and triple-loop learning initiatives might be beneficial if objectives and assumptions need reframing or if the underlying beliefs and values have transformed (Pahl-Wostl 2009), especially in a changing environmental and social climate. Maintaining all eyes on the end objective (i.e., the prize) of healthy sustainable Great Lakes fisheries requires dedicated communications and outreach efforts across multiple jurisdictions and publics and carrying forward institutional knowledge across generations of Great Lakes stewards.

Conclusions

The 10 lessons presented in the present article (figure 4) represent ones that have staying power in that they have been realized repeatedly or continuously over at least the last three decades. There are more lessons that we opted to omit because they were specific to a given project or control measure but less germane to the control program as a whole. We contend that the lessons shared in the present article, in whole or in part, can be applied in diverse contexts that extend well beyond aquatic ecosystems or North America. It is important to emphasize that, since its inception, the control program has been supported by a parallel mission-oriented science program to support management. Recognizing that science can exist in a vacuum, the GLFC has also embraced approaches to connect managers with scientists by embracing coproduction when possible, including managers on grant-making committees, and implementing a science transfer



Figure 4. Summary of the 10 lessons for controlling invasive sea lamprey that are germane to other invasive species control programs. The control program is supported by a science program (which funds research on lamprey biology and control) with a science transfer program serving a key role in knowledge exchange and translation. The GLFC is in the early days of its journey to create space for Indigenous knowledge systems and ways of knowing in the sea lamprey control program.

program to facilitate knowledge exchange among researchers, decision-makers, and sea lamprey control agents (Hinderer et al. 2021). The GLFC has long embraced a framework that uses military strategy and verbiage (and is not alone among invasive species control programs; Janovsky and Larson 2019), extending back to Sun Tzu's centuries-old *The Art of War* (see Tzu 2008) and which extends across the lessons (e.g., know your enemy). Consider thinking about the art of invasive species control. Those historic teachings extend from laying plans to waging war to attack by stratagem. Notably, Tzu stated that "If you know the enemy and know yourself, you need not fear the result of a hundred battles. If you know yourself but not the enemy, for every victory gained you will also suffer a defeat. If you know neither the enemy nor yourself, you will succumb in every battle." In the sea lamprey context, that means know your strengths and weaknesses (both internally and with respect to the target invasive species) and allocate available but limited resources where you can have the most positive influence on the ecosystem (e.g., minimizing the cost per kill or exploiting stereotyped behavioral responses of sea lamprey to traps). We acknowledge that making knowing your enemy a focal point is a challenging task. The GLFC and allied agencies had focused on both knowing their enemy (using science and adaptive management and monitoring) but have also taken the time to understand and build their own capacity and capabilities (recognizing an evolution in thinking). When, at times, actions failed, those failures were not simply dismissed but rather dissected to learn and improve (e.g., Marcy-Quay et al. 2025). We acknowledge that what an invasive species is continues to evolve (recently informed by Indigenous learnings; see Black et al. 2021, Mattes and Kitson 2021, Wehi et al. 2023) such that the military framing may eventually become inappropriate or irrelevant. Whether the control program philosophy is rooted in centuries old military stratagem or a more modern philosophy of caring and sharing, a dedicated control program with dedicated people is critical. To achieve success, such a program needs to be broadly supported, well resourced, coordinated across jurisdictions and integrated across control program elements, knowledge based, diverse, flexible and adaptive, and determined.

Most invasive species control programs do not have the history or resources (control agents, funding, and science) that exist for

sea lamprey in the Great Lakes basin. That is more reason to look toward established programs and institutions such as the control program and GLFC for guidance. Most of the lessons shared in the present article can be downscaled or adapted to a more localized or nascent context in that planning for invasive species control is a long-term commitment. As outlined in the present article, through approaches such as adaptive management, horizon scanning methods, and embracing implementation science, the lessons learned about sea lamprey control will continue to evolve, which is itself a lesson. A responsibility to do better and the institutional culture to do so are critical for ensuring that evidence and not emotion guide invasive species control programs. We submit that the lessons shared in the present article will help to guide invasive species control programs spanning taxa, ecosystems, and regions.

Acknowledgments

Tyler Buchinger, Anne Scott, and Robert McLaughlin provided comments on an earlier version of the manuscript. We thank the members of the Great Lakes Fishery Commission Sea Lamprey Research Board and the Sea Lamprey Control Program for their inspiration for this project. We thank Kathryn Peiman for formatting the references. Any use of trade, product, or firm name is for descriptive purposes only and does not imply endorsement by the US government. We thank several anonymous referees for providing thoughtful comments on the manuscript.

Author Contributions

Steven J Cooke (Conceptualization, Investigation, Project administration, Visualization, Writing – original draft, Writing – review & editing), Carrie L. Baker (Conceptualization, Writing – original draft, Writing – review & editing), Julie L. Mida Hinderer (Conceptualization, Writing – original draft, Writing – review & editing), Michael Siefkes (Writing – original draft, Writing – review & editing), Jessica M. Barber (Writing – original draft, Writing – review & editing), Todd B. Steeves (Writing – original draft, Writing – review & editing), Margaret Docker (Conceptualization, Writing – original draft, Writing – review & editing), Weiming Li (Conceptualization,

Writing – original draft, Writing – review & editing), Michael P Wilkie (Conceptualization, Writing – original draft, Writing – review & editing), Michael L. Jones (Conceptualization, Writing – original draft, Writing – review & editing), Erin Dunlop (Conceptualization, Writing – original draft, Writing – review & editing), Cory O. Brant (Writing – original draft, Writing – review & editing), Nicholas Johnson (Conceptualization, Writing – original draft, Writing – review & editing), William P. Mattes (Writing – original draft, Writing – review & editing), Marc Gaden (Conceptualization, Writing – original draft, Writing – review & editing), Andrew Muir (Conceptualization, Project administration, Resources, Visualization, Writing – original draft, Writing – review & editing).

Funding was provided by the Great Lakes Fishery Commission. SJC, MFD, and MPW are also supported by the NSERC Discovery Grant program and SJC and MFD are also part of the Genome Canada project GenFish.

References cited

- Ahmed DA, et al. 2023. Recent advances in availability and synthesis of the economic costs of biological invasions. *BioScience* 73: 560–574.
- Allen CR, Gunderson LH. 2011. Pathology and failure in the design and implementation of adaptive management. *Journal of Environmental Management* 92: 1379–1384.
- Almack K, Dunlop ES, Lauzon R, Nadjiwon S, Duncan AT. 2023. Building trust through the two-eyed seeing approach to joint fisheries research. *Journal of Great Lakes Research* 49: S46–S57.
- Bartlett C, Marshall M, Marshall A. 2012. Two-eyed seeing and other lessons learned within a co-learning journey of bringing together indigenous and mainstream knowledges and ways of knowing. *Journal of Environmental Studies and Sciences* 2: 331–340.
- Behnken JA, Groninger JW, Akamani K. 2016. Institutional constraints to collaborative ecosystem management within a wetlands conservation partnership. *Journal of Contemporary Water Research and Education* 158: 19–33.
- Bellard C, Genovesi P, Jeschke JM. 2016. Global patterns in threats to vertebrates by biological invasions. *Proceedings of the Royal Society B* 283: 20152454.
- Bellard C, Bernery C, Leclerc C. 2021. Looming extinctions due to invasive species: Irreversible loss of ecological strategy and evolutionary history. *Global Change Biology* 27: 4967–4979.
- Black A, Garner G, Mark-Shadbolt M, Balanovic J, MacDonald E, Mercier O, Wright J. 2021. Indigenous peoples' attitudes and social acceptability of invasive species control in New Zealand. *Pacific Conservation Biology* 28: 481–490.
- Bomford M, O'Brien P. 1995. Eradication or control for vertebrate pests? *Wildlife Society Bulletin* 1995: 249–255.
- Brant C. 2019. *Great Lakes Sea Lamprey: The 70 Year War on a Biological Invader*. University of Michigan Press.
- Brant CO, Read J, Gaden M, Shaw R. 2019. *Great Lakes: Lessons from Aquatic Invasive Species Cases to Inform Future Invasive Species Policy and Practice*. University of Michigan Water Center. <https://graham.umich.edu/media/pubs/Great-Lakes-AIS-Report-Case-Study-45903.pdf>
- Brisaki E, et al. 2013. Multidimensional approach to invasive species prevention. *Environmental Science and Technology* 47: 1216–1221.
- Buchinger TJ, Siefkes MJ, Zielinski BS, Brant CO, Li W. 2015. Chemical cues and pheromones in the sea lamprey (*Petromyzon marinus*). *Frontiers in Zoology* 12: 32.
- Burgiel SW. 2020. The incident command system: A framework for rapid response to biological invasion. *Biological Invasions* 22: 155–165.
- Burkett DP, Barber JM, Steeves TB, Siefkes MJ. 2021. Sea lamprey control 2020–2040: Charting a course through dynamic waters. *Journal of Great Lakes Research* 47: S809–S814.
- Carneiro L, et al. 2024. Benefits do not balance costs of biological invasions. *BioScience* 74: 340–344.
- Cheruvellil KS, Soranno PA, Weathers KC, Hanson PC, Goring SJ, Filstrup CT, Read EK. 2014. Creating and maintaining high-performing collaborative research teams: The importance of diversity and interpersonal skills. *Frontiers in Ecology and the Environment* 12: 31–38.
- Coleman MJ, Sindel BM, Stayner RA. 2017. Effectiveness of best practice management guides for improving invasive species management: A review. *Rangeland Journal* 39: 39–48.
- Cooke SJ, Young N, Alexander S, Kadykalo AN, Danylchuk AJ, Muir AM, Hinderer JLM, Cvitanovic C, Nguyen VM. 2024. Embracing implementation science to enhance fisheries and aquatic management and conservation. *Fisheries* 49: 475–485.
- Comicelli L, Whitehead J, Bragg L, Lange TZ, Southwick R. 2022. *Economic Aspects of the Great Lakes Recreational Fisheries and Factors Driving Change*. Great Lakes Fishery Commission. Project completion report, project ID=2020_ALL_440910. https://www.glf.org/pubs/pdfs/research/reports/2020_ALL_440910.pdf
- Crystal-Ornelas R, Lockwood JL. 2020. The 'known unknowns' of invasive species impact measurement. *Biological Invasions* 22: 1513–1525.
- Daghfous G, Green WW, Alford ST, Zielinski BS, Dubuc R. 2016. Sensory activation of command cells for locomotion and modulatory mechanisms: Lessons from lampreys. *Frontiers in Neural Circuits* 10: 18.
- Dahlberg AD, Waller DL, Hammond D, Lund K, Phelps NBD. 2023. Open water dreissenid mussel control projects: Lessons learned from a retrospective analysis. *Scientific Reports* 13: 10410
- Davies SJ, et al. 2020. Experience and lessons from alien and invasive animal control projects in South Africa. Pages 629–663 in van Wilgen BW, Measey J, Richardson DM, Wilson JR, Zengeya TA, eds. *Biological Invasions in South Africa*. Springer.
- Diagne C, Leroy B, Vaissière AC, Gozlan RE, Roiz D, Jarić I, Salles J-M, Bradshaw CJA, Courchamp F. 2021. High and rising economic costs of biological invasions worldwide. *Nature* 592: 571–576.
- Duncan AT, Lauzon R, Harpur C. 2023. An investigation into Saugeen Ojibway Nation-based ecological knowledge on the ciscoes (*Coregonus* spp.) of Lake Huron. *Journal of Great Lakes Research* 49: S138–S147.
- Dunlop ES, et al. 2018. Rapid evolution meets invasive species control: The potential for pesticide resistance in sea lamprey. *Canadian Journal of Fisheries and Aquatic Sciences* 75: 152–168.
- Eade D. 2007. Capacity building: Who builds whose capacity? *Development in Practice* 17: 630–639.
- Evans TM, Wagner CM, Miehl SM, Johnson NS, Haas TF, Dunlop E, Manzon RG. 2021. Before the first meal: The elusive pre-feeding juvenile stage of the sea lamprey. *Journal of Great Lakes Research* 47: S580–S589.
- Ferreira-Martins D, Champer J, McCauley DW, Zhang Z, Docker MF. 2021. Genetic control of invasive sea lamprey in the Great Lakes. *Journal of Great Lakes Research* 47: S764–S775.
- Gaden M, Brant CO, Lambe R. 2021. Why a Great Lakes Fishery Commission? The seven-decade pursuit of a Canada-US fishery treaty. *Journal of Great Lakes Research* 47: S11–S23.
- Gaden M, Brant CO, Stedman RC, Cooke SJ, Young N, Lauber TB, Nguyen VM, Connolly NA, Knuth B. 2021. Shifting baselines

- and social license to operate: Challenges in communicating sea lamprey control. *Journal of Great Lakes Research* 47: S800–S808.
- Gaden M, Brant CO, Lambe R. 2022. The History behind the Canada–U.S. Convention on Great Lakes Fisheries: A Seven-Decade Effort. *Great Lakes Fishery Commission. Laurentian report no. 2022-01*
- Haack RA, Hérard F, Sun J, Turgeon JJ. 2010. Managing invasive populations of Asian longhorned beetle and citrus longhorned beetle: A worldwide perspective. *Annual Review of Entomology* 55: 521–546.
- Haley AL, et al. 2023. On the effectiveness of public awareness campaigns for the management of invasive species. *Environmental Conservation* 50: 202–211.
- Hall TC. 1996. Operational control of the brown tree snake on Guam. Pages 234–240 in Crabb AC, eds. *Proceedings of the Vertebrate Pest Conference*. University of California Davis.
- Hansen GJA, Jones ML. 2008. A rapid assessment approach to prioritizing streams for control of Great Lakes sea lampreys: A case study in adaptive management. *Canadian Journal of Fisheries and Aquatic Sciences* 65: 2471–2484.
- Hansen GJA, Jones ML. 2008. The value of information in fishery management. *Fisheries* 33: 340–348.
- Herbst SJ, et al. 2021. An adaptive management approach for implementing multi-jurisdictional response to grass carp in Lake Erie. *Journal of Great Lakes Research* 47: 96–107.
- Hinderer JLM, et al. 2021. Insights from a novel, user-driven science transfer program for resource management. *Socio-Ecological Practice Research* 3: 337–362.
- Holmes ND, Griffiths R, Pott M, Alifano A, Will D, Wegmann AS, Russell JC. 2015. Factors associated with rodent eradication failure. *Biological Conservation* 185: 8–16.
- Humphries T, Campbell S, Florentine S. 2022. Challenges inherent in controlling prickly pear species; a global review of the properties of *Opuntia stricta*, *Opuntia ficus-indica* and *Opuntia monacantha*. *Plants* 11: 3334.
- [IPBES] Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. 2019. *Global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services*. IPBES. <https://doi.org/10.5281/zenodo.3553579>.
- Janovsky RM, Larson ER. 2019. Does invasive species research use more militaristic language than other ecology and conservation biology literature? *NeoBiota* 44: 27–38.
- Jardine SL, Sanchirico JN. 2018. Estimating the cost of invasive species control. *Journal of Environmental Economics and Management* 87: 242–257.
- Jellinek S, Llyod S, Catterall C, Sato C. 2021. Facilitating collaboration between researchers and practitioners in ecosystem management and restoration. *Ecological Management and Restoration* 22: 208–213.
- Johnson FA, Smith BJ, Bonneau M, Martin J, Romagosa C, Mazzotti F, Waddle H, Reed RN, Eckles JK, Vitt LJ. 2017. Expert elicitation, uncertainty, and the value of information in controlling invasive species. *Ecological Economics* 137: 83–90.
- Johnson NS, Lewandoski SA, Jubar AK, Symbal MJ, Solomon BM, Bravener GA, Barber JM, Siefkes MJ. 2024. A decade-long study demonstrates that a population of invasive sea lamprey (*Petromyzon marinus*) can be controlled by introducing sterilized males. *Scientific Reports* 14: 12689.
- Jones ML, Irwin B, Hansen GJA, Dawson HA, Treble AJ, Liu W, Dai W, Bence JR. 2009. An operating model for Great Lakes sea lamprey integrated pest management. *Open Fish Science Journal* 2: 59–73.
- Jubar AK, Frank RJ, Keffer DA, Neave FB, Symbal MJ, Steeves TB. 2021. Prioritizing lampricide treatments in Great Lakes tributaries and lentic areas: An adaptive management approach to sea lamprey control during 2000–2017. *Journal of Great Lakes Research* 47: S238–S246.
- Keller RP, Frang K, Lodge DM. 2008. Preventing the spread of invasive species: Economic benefits of intervention guided by ecological predictions. *Conservation Biology* 22: 80–88.
- Koel TM, et al. 2020. Yellowstone Lake ecosystem restoration: A case study for invasive fish management. *Fishes* 5: 18.
- Krueger CC, Marsden JE. 2007. Sea lamprey research: Balancing basic and applied research, and using themes to define priorities. *Journal of Great Lakes Research* 33: 1–6.
- Labin SN, Duffy JL, Meyers DC, Wandersman A, Lesesne CA. 2012. A research synthesis of the evaluation capacity building literature. *American Journal of Evaluation* 33: 307–338.
- Leung B, Lodge DM, Finnoff D, Shogren JF, Lewis MA, Lamberti G. 2002. An ounce of prevention or a pound of cure: Bioeconomic risk analysis of invasive species. *Proceedings of the Royal Society B* 269: 2407–2413.
- Lewandoski SA, Brenden TO, Siefkes MJ, Johnson NS. 2021. An adaptive management implementation framework for evaluating supplemental sea lamprey (*Petromyzon marinus*) controls in the Laurentian Great Lakes. *Journal of Great Lakes Research* 47: S753–S763.
- Lewandoski SA, et al. 2025. Decision analysis of integrated pest management: A case study on invasive sea lamprey in the Great Lakes Basin. *Journal of Environmental Management* 373: 123666.
- Li K, et al. 2018. Fatty-acid derivative acts as a sea lamprey migratory pheromone. *Proceedings of the National Academy of Sciences* 115: 8603–8608.
- Li W, Sorensen PW, Gallaher DD. 1995. The olfactory system of migratory adult sea lamprey (*Petromyzon marinus*) is specifically and acutely sensitive to unique bile acids released by conspecific larvae. *Journal of General Physiology* 105: 569–587.
- Lieurance D, et al. 2023. Identifying invasive species threats, pathways, and impacts to improve biosecurity. *Ecosphere* 14: e4711.
- Linders TE, et al. 2020. The impact of invasive species on social-ecological systems: Relating supply and use of selected provisioning ecosystem services. *Ecosystem Services* 41: 101055.
- Mack RN, Simberloff D, Mark Lonsdale W, Evans H, Clout M, Bazzaz FA. 2000. Biotic invasions: Causes, epidemiology, global consequences, and control. *Ecological Applications* 10: 689–710.
- Marcy-Quay B, et al. 2025. Sea lamprey control reduction during the COVID-19 pandemic corresponds to rapid increase in Sea Lamprey abundance. *Fisheries* 2025: vua020.
- Mattes WP, Kitson JC. 2021. Sea lamprey control in the Great Lakes: A tribal/First Nations representative's perspective. *Journal of Great Lakes Research* 47: S796–S799.
- McCauley DW, Docker MF, Whyard S, Li W. 2015. Lampreys as diverse model organisms in the genomics era. *BioScience* 65: 1046–1056.
- McGeoch MA, Butchart SH, Spear D, Marais E, Kleynhans EJ, Symes A, Chanson J, Hoffmann M. 2010. Global indicators of biological invasion: Species numbers, biodiversity impact and policy responses. *Diversity and Distributions* 16: 95–108.
- McLaughlin R, et al. 2021. Foreword: Control and conservation of lampreys beyond 2020—Proceedings from the 3rd Sea Lamprey International Symposium (SLIS III). *Journal of Great Lakes Research* 47: S1–S10.
- Mehta SV, Haight RG, Homans FR, Polasky S, Venette RC. 2007. Optimal detection and control strategies for invasive species management. *Ecological Economics* 61: 237–245.
- Miehls S, Sullivan P, Twohey M, Barber J, McDonald R. 2020. The future of barriers and trapping methods in the sea lamprey

- (*Petromyzon marinus*) control program in the Laurentian Great Lakes. *Reviews in Fish Biology and Fisheries* 30: 1–24.
- Muir AM, Krueger CC, Hansen MJ. 2012. Re-establishing lake trout in the Laurentian Great Lakes: Past, present, and future. Pages 533–588 in Taylor WW, Lynch AJ, Leonard NJ, eds. *Great Lakes Fisheries Policy and Management: A Binational Perspective*, 2nd ed. Michigan State University Press.
- Muir AM, et al. 2023. Confronting a post-pandemic new normal: Threats and opportunities to trust-based relationships in natural resource science and management. *Journal of Environmental Management* 330: 117140.
- Mullin AE, Coe IR, Gooden EA, Tunde-Byass M, Wiley RE. 2021. Inclusion, diversity, equity, and accessibility: From organizational responsibility to leadership competency. *Healthcare Management Forum* 34: 311–315.
- Mulvaney KK, Lee S, Höök TO, Prokopy LS. 2015. Casting a net to better understand fisheries management: An affiliation network analysis of the Great Lakes Fishery Commission. *Marine Policy* 57: 120–131.
- Myers JH, Simberloff D, Kuris AM, Carey JR. 2000. Eradication revisited: Dealing with exotic species. *Trends in Ecology and Evolution* 15: 316–320.
- Nguyen VM, Young N, Cooke SJ. 2017. A roadmap for knowledge exchange and mobilization research in conservation and natural resource management. *Conservation Biology* 31: 789–798.
- Nonkes C, Duncan AT, Lauzon R, Ryan K, Reid AJ, Cooke SJ, Young N. 2023. Two-eyed seeing: Developing perspective and wisdom on sea lamprey in the Laurentian Great Lakes. *Journal of Great Lakes Research* 49: S148–S159.
- Nonkes C, Lauzon R, Redford B, Duncan AT. 2024. Adaptation through knowledge coexistence: Insights for environmental and sea lamprey stewardship. *Local Environment* 29: 951–968.
- Pahl-Wostl C. 2009. A conceptual framework for analysing adaptive capacity and multi-level learning processes in resource governance regimes. *Global Environmental Change* 19: 354–365.
- Phillips C, Brown K, Broome K, Green C, Walker G. 2019. Criteria to help evaluate and guide attempts to eradicate arthropod pests. Pages 400–404 in Veitch CR, Clout MN, Martin AR, Russell JC, West CJ eds. *Island Invasives: Scaling Up to Meet the Challenge*. International Union for Conservation of Nature. Occasional Paper SSC no. 62.
- Rai PK, Singh JS. 2020. Invasive alien plant species: Their impact on environment, ecosystem services, and human health. *Ecological Indicators* 111: 106200.
- Reaser JK, Burgiel SW, Kirkey J, Brantley KA, Veatch SD, Burgos-Rodríguez J. 2020. The early detection of and rapid response (EDRR) to invasive species: A conceptual framework and federal capacities assessment. *Biological Invasions* 22: 1–19.
- Reid AJ, Eckert LE, Lane JF, Young N, Hinch SG, Darimont CT, Cooke SJ, Ban NC, Marshall A. 2021. “Two-eyed seeing”: An Indigenous framework to transform fisheries research and management. *Fish and Fisheries* 22: 243–261.
- Robinson KF, Miehl SM, Siefkes MJ. 2021a. Understanding sea lamprey populations in the Great Lakes prior to broad implementation of sea lamprey control. *Journal of Great Lakes Research* 47: S328–S334.
- Robinson KF, et al. 2021. Using decision analysis to collaboratively respond to invasive species threats: A case study of Lake Erie grass carp (*Ctenopharyngodon idella*). *Journal of Great Lakes Research* 47: 108–119.
- Samaniego A, et al. 2021. Factors leading to successful island rodent eradications following initial failure. *Conservation Science and Practice* 3: e404.
- Sawyer AJ. 1980. Prospects for integrated pest management of the sea lamprey (*Petromyzon marinus*). *Canadian Journal of Fisheries and Aquatic Sciences* 37: 2081–2092.
- Scott AM, Johnson NS, Siefkes MJ, Li W. 2023. Synergistic behavioral antagonists of a sex pheromone reduce reproduction of invasive sea lamprey. *iScience* 26: 107744.
- Shanmuganathan T, et al. 2010. Biological control of the cane toad in Australia: A review. *Animal Conservation* 13: 16–23.
- Siefkes MJ. 2017. Use of physiological knowledge to control the invasive sea lamprey (*Petromyzon marinus*) in the Laurentian Great Lakes. *Conservation Physiology* 5: cox031.
- Siefkes MJ, Johnson NS, Muir AM. 2021. A renewed philosophy about supplemental sea lamprey controls. *Journal of Great Lakes Research* 47: S742–S752.
- Simberloff D. 2009. We can eliminate invasions or live with them. Successful management projects. *Biological Invasions* 11: 149–157.
- Simberloff D. 2014. Biological invasions: What’s worth fighting and what can be won? *Ecological Engineering* 65: 112–121.
- Smith SH. 1968. Species succession and fishery exploitation in the Great Lakes. *Journal of the Fisheries Research Board of Canada* 25: 667–693.
- Smith JJ, Kuraku S, Holt C, Sauka-Spengler T, Jiang N, Campbell MS, Yandell MD, Manousaki T, Meyer A, Bloom OE. 2013. Sequencing of the sea lamprey (*Petromyzon marinus*) genome provides insights into vertebrate evolution. *Nature Genetics* 45: 415–421.
- Smith JJ, Timoshevskaya N, Ye C, Holt C, Keinath MC, Parker HJ, Cook ME, Hess JE, Narum SR, Lamanna F. 2018. The sea lamprey germline genome provides insights into programmed genome rearrangement and vertebrate evolution. *Nature Genetics* 50: 270–277.
- Sobeck J, Agius E. 2007. Organizational capacity building: Addressing a research and practice gap. *Evaluation and Program Planning* 30: 237–246.
- Song AM, Temby O, Kim D, Saavedra Cisneros A, Hickey GM. 2019. Measuring, mapping and quantifying the effects of trust and informal communication on transboundary collaboration in the Great Lakes fisheries policy network. *Global Environmental Change* 54: 6–18.
- Song AM, Temby O, Kim D, Hickey GM. 2020. Assessing the influence of international environmental treaty secretariats using a relational network approach. *Earth System Governance* 5: 100076.
- Sower SA. 2015. The reproductive hypothalamic–pituitary axis in lampreys. Pages 305–373 in Docker MF, ed. *Lampreys: Biology, Conservation, and Control*. Springer.
- Sower SA. 2018. Landmark discoveries in elucidating the origins of the hypothalamic–pituitary system from the perspective of a basal vertebrate, sea lamprey. *General and Comparative Endocrinology* 264: 3–15.
- Sullivan WP, et al. 2021. The path toward consistent achievement of sea lamprey abundance and lake trout marking targets in Lake Ontario, 2000–2019. *Journal of Great Lakes Research* 47: S523–S548.
- Swink WD. 2003. Host selection and lethality of attacks by sea lampreys (*Petromyzon marinus*) in laboratory studies. *Journal of Great Lakes Research* 29: 307–319.
- Tzu S. 2008. The art of war. Pages 63–91 in Mahnken TG, Maiolo JA, eds. *Strategic Studies*. Routledge.
- van Putten I, Ison S, Cvitanovic C, Hobday AJ, Thomas L. 2022. Who has influence? The role of trust and communication in the conservation of flatback turtles in Western Australia. *Regional Studies in Marine Science* 49: 102080.
- Vilà M, Hulme PE. 2017. *Impact of Biological Invasions on Ecosystem Services*, vol. 12. Springer.

- Walters CJ. 1986. *Adaptive Management of Renewable Resources*. Basingstoke, UK: MacMillan Publishing Company.
- Wehi PM, Kamelamela KL, Whyte K, Watene K, Reo N. 2023. Contribution of Indigenous peoples' understandings and relational frameworks to invasive alien species management. *People and Nature* 5: 1403–1414.
- Wiener CS, Rivera MA, Toonen RJ, Leong JAC, Kosaki RK, Karl S, Keller K, Johnson H. 2011. Creating effective partnerships in ecosystem-based management: A culture of science and management. *Journal of Marine Science* 2011: 241610.
- Wilkie MP, Hubert TD, Boogaard MA, Birceanu O. 2019. Control of invasive sea lampreys using the piscicides TFM and niclosamide: Toxicology, successes and future prospects. *Aquatic Toxicology* 211: 235–252.
- Wingfield J, Brant C, Eshenroder R, Gaden M, Miehl A, Siefkes M. 2021. 100 years of sea lampreys above Niagara Falls: A reflection on what happened and what we learned. *Journal of Great Lakes Research* 47: 1844–1848.
- Wylie FR, Janssen-May S. 2017. Red imported fire ant in Australia: What if we lose the war? *Ecological Management and Restoration* 18: 32–44.
- York JR, Lee EM, McCauley DW. 2019. The lamprey as a model vertebrate in evolutionary developmental biology. Pages 481–526 in Docker MF, ed. *Lampreys: Biology, Conservation, and Control*. Springer.
- Zhang Z, Zhang Q, Dexheimer TS, Ren J, Neubig RR, Li W. 2020. Two highly related odorant receptors specifically detect α -bile acid pheromones in sea lamprey (*Petromyzon marinus*). *Journal of Biological Chemistry* 295: 12153–12166.
- Zielinski DP, McLaughlin R, Castro-Santos T, Paudel B, Hrodey P, Muir A. 2019. Alternative sea lamprey barrier technologies: History as a control tool. *Reviews in Fish Science and Aquaculture* 27: 438–457.
- Zielinski DP, McLaughlin RL, Pratt TC, Goodwin RA, Muir AM. 2020. Single-stream recycling inspires selective fish passage solutions for the connectivity conundrum in aquatic ecosystems. *BioScience* 70: 871–886.

Received: March 21, 2025. Revised: July 16, 2025. Accepted: July 28, 2025

© The Author(s) 2025. Published by Oxford University Press on behalf of the American Institute of Biological Sciences. This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact reprints@oup.com for reprints and translation rights for reprints. All other permissions can be obtained through our RightsLink service via the Permissions link on the article page on our site—for further information please contact journals.permissions@oup.com