

THE EFFECT OF NATURAL DISSOLVED ORGANIC CARBON ON THE ACUTE TOXICITY OF COPPER TO LARVAL FRESHWATER MUSSELS (GLOCHIDIA)

PATRICIA L. GILLIS,*†‡ JAMES C. MCGEER,§ GERALD L. MACKIE,† MICHAEL P. WILKIE,§ and JOSEF D. ACKERMAN†

†Department of Integrative Biology, University of Guelph, Guelph, Ontario N1G-2W1, Canada

‡Aquatic Ecosystem Protection Research Division, Environment Canada, Burlington, Ontario L7R-4A6, Canada

§Department of Biology, Wilfrid Laurier University, Waterloo, Ontario N2L-3C5, Canada

(Submitted 28 September 2009; Returned for Revision 23 December 2009; Accepted 9 June 2010)

Abstract—The present study examined the effect of dissolved organic carbon (DOC), both added and inherent, on Cu toxicity in glochidia, the larvae of freshwater mussels. Using incremental additions of natural DOC concentrate and reconstituted water, a series of acute copper toxicity tests were conducted. An increase in DOC from 0.7 to 4.4 mg C/L resulted in a fourfold increase (36–150 µg Cu/L) in the 24-h median effective concentration (EC50) and a significant linear relationship ($r^2 = 0.98$, $p = 0.0008$) between the DOC concentration and the Cu EC50 of *Lampsilis siliquoidea* glochidia. The ameliorating effect of added DOC on Cu toxicity was confirmed using a second mussel species, the endangered (in Canada) *Lampsilis fasciola*. The effect of inherent (i.e., not added) DOC on Cu toxicity was also assessed in eight natural waters (DOC 5–15 mg C/L). These experiments revealed a significant relationship between the EC50 and the concentration of inherent DOC ($r^2 = 0.79$, $p = 0.0031$) with EC50s ranging from 27 to 111 µg Cu/L. These laboratory tests have demonstrated that DOC provides glochidia with significant protection from acute Cu toxicity. The potential risk that Cu poses to mussel populations was assessed by comparing Cu and DOC concentrations from significant mussel habitats in Ontario to the EC50s. Although overall mean Cu concentration in the mussel's habitat was well below the acutely toxic level given the concentration of DOC, episodic Cu releases in low DOC waters may be a concern for the recovery of endangered freshwater mussels. The results are examined in the context of current Cu water quality regulations including the U.S. Environmental Protection Agency's (U.S. EPA) biotic ligand model. Environ. Toxicol. Chem. 2010;29:2519–2528. © 2010 SETAC

Keywords—Copper toxicity Dissolved organic carbon Freshwater mussels Glochidia Biotic ligand model

INTRODUCTION

Freshwater mollusks are among the most endangered groups of organisms in North America, with nearly 70% of the species designated as either threatened, endangered, or in decline [1]. Their decline has been attributed to a number of factors, including habitat alteration, loss of fish hosts, invasive species, and exposure to environmental pollution [2–5]. Some studies [2,6] have suggested that environmental contamination may contribute to recruitment or reproductive failure in freshwater mussels. As with most organisms, the early life stages of freshwater mussels are the most sensitive to environmental pollution. The larvae, called glochidia, are obligate parasites on fish. Most species of freshwater mussels release their glochidia from the brooding chambers in the gills of the female into the water column, where on contact they will encyst upon a host. The amount of time that a given glochidium is in the water column and exposed to waterborne contaminants can vary significantly (seconds to weeks) depending on the specific reproductive strategies of the species (luring or broadcasting) and how long it takes to make contact with a host [7]. Although freshwater mussels are sensitive to a range of pollutants, they are particularly sensitive to Cu [6,8–11]. In fact, concerns have been expressed that water quality guidelines for Cu do not adequately protect the early life stages of freshwater mussels [12,13]. It is important that toxicity data for this sensitive early

life stage be included when water quality regulations are developed, because contaminants that reduce glochidia success will put mussel populations at risk.

As with all metals, Cu bioavailability is controlled by the composition of the exposure water. Major ions, including Ca, Mg, and Na, compete with metal ions at the site of uptake [14,15]. Similarly, the presence of ligands, such as dissolved organic carbon (DOC), can bind with the metal ion and reduce its bioavailability [16]. Copper bioavailability is strongly influenced by DOC. Free Cu ions (Cu^{2+}), the most bioavailable and thus toxic Cu species in solution [17], form complexes with the negatively charged natural organic matter, reducing the amount of free Cu in solution and thereby reducing the bioavailability of Cu [18]. In fact, Mantoura et al. [19] reported that greater than 90% of Cu in freshwater was complexed by humic materials. The effect of copper-DOC complexation on the bioavailability and toxicity of Cu has been demonstrated in a range of aquatic organisms, from cladocerans to fish [20–25], but until recently information on the effect of DOC on Cu toxicity in freshwater mussels was relatively scarce. Wang et al. [26] evaluated the effect of natural DOC on acute Cu toxicity in juvenile freshwater mussels. They observed a significant linear increase in the Cu EC50 (7–12-fold) when exposure solutions were augmented with up to 10 mg/L of natural DOC. Markich et al. [27] investigated the effect of DOC (as fulvic acid) on the response of adult *Hyridella depressa* (Australian freshwater river mussel) to metals. They reported an 18-fold decrease in Cu sensitivity (as measured in terms of duration of valve opening) when the concentration of fulvic acid in the exposures was increased from 0 to 11 mg/L. Hanstén et al. [28] reported that additions of

* To whom correspondence may be addressed
(patty.gillis@ec.gc.ca).

Published online 9 July 2010 in Wiley Online Library
(wileyonlinelibrary.com).

humic acid (color value of 200 mg Pt/L) reduced the toxicity of Cd and Cu to *Anodonta anatine* (duck mussel) glochidia. Also with glochidia, small additions (0.3–0.6 mg C/L) of a standard humic substance (Aldrich humic acid) resulted in significant decreases in Cu toxicity of *Epioblasma triquetra* (snuffbox) and *Lampsilis fasciola* (wavy-rayed lampmussel) glochidia [11]. However, it is still unknown how effective natural DOC from aquatic systems would be at protecting larval freshwater mussels from acute Cu exposure.

Given the significant relationship between DOC and Cu bioavailability, along with the heightened sensitivity of freshwater mussels to Cu, it is important to understand the influence of DOC on glochidia Cu sensitivity in order to assess the potential threat that Cu exposure could have on the recovery of imperiled freshwater mussels. The present study, therefore, examines the effect of natural DOC, both added and inherent (i.e., not added) on the Cu toxicity of the larvae of two species of freshwater mussels. Specifically, the influence of DOC on acute Cu toxicity of glochidia was examined using a reconstituted soft water and incremental additions of a natural organic matter, and also using a selection of source waters with their inherent natural organic matter (DOC varied from 5–15 mg C/L). These data are discussed in relation to current water quality criteria as well as the implications for populations of freshwater mussels.

MATERIALS AND METHODS

Mussel collection and laboratory care

Gravid *Lampsilis fasciola* (Rafinesque 1820) and *Lampsilis siliquoidea* (Barnes 1823) (fatmucket) were collected from the Thames River, and the latter were also collected Cox Creek (Grand River Watershed), both in Ontario, Canada. *L. fasciola*, which has been designated as an endangered species in Canada by the Committee on the Status of Endangered Wildlife in Canada (COSEWIC), was collected under the Species at Risk Permit Number SECT 06 SCI 007. The mussels were maintained in the Aqualab facility at the University of Guelph, (Guelph, ON, Canada) in a flow-through system using well water at $10 \pm 2^\circ\text{C}$ to prevent the release of glochidia. The mussels were fed a commercial shellfish diet (Instant Algae Shellfish Diet 1800[®]) which is a mixture of a *Nannochloropsis*, *Isochrysis*, *Pavlova*, *Thalassiosira*, and *Tetraselmis* algae at a rate of approximately 1.2×10^{10} algae cells per mussel per day.

Glochidia were collected by flushing the marsupia (brooding chambers) with water using a syringe. The viability of each mussel's glochidia was assessed prior to use (viability procedure described below). For each exposure, glochidia were collected from a minimum of three gravid females that exhibited greater than 90% viability. Prior to initiating an exposure, the pooled glochidia sample was gradually (over a 2-h period) brought up to the exposure temperature of 21°C by dilutions with room temperature reconstituted water.

Acute copper toxicity tests with glochidia

Toxicity tests were modeled after the American Society for Testing and Materials method [29] for conducting toxicity tests with the early life stages of freshwater mussels. Briefly, glochidia were removed from gravid female mussels and exposed to various concentrations of a waterborne contaminant. After a given period of exposure (24 h), the viability of the exposed glochidia (i.e., ability to close their valves and clamp down on a fish's gill for encystment) was assessed using a concentrated salt solution (NaCl 240 g/L). Glochidia viability was assessed in a subsample (100 to 200) of the approximately 1,000 glochidia

exposed in each replicate. Viability was calculated using the following equation: Percent Viability = $100 \times (\text{Number of closed glochidia after addition of NaCl} - \text{Number of closed glochidia before addition of NaCl}) / (\text{Number of closed glochidia after addition of NaCl} + \text{Number of open glochidia after addition of NaCl})$. Results are expressed as EC50 values rather than median lethal concentration (LC50) values, but because they are obligatory parasites on fish, for practical purposes nonviable glochidia should be considered dead, because they would be unable to attach to a fish host and complete their life cycle.

An aqueous Cu stock solution made from certified American Chemical Society grade (Fisher Scientific) cupric sulfate ($\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$) was used to create the copper exposures. Copper spiked-DOC exposure solutions (described below) were acclimated to the test temperature (21°C) before addition of glochidia. Exposures (100 ml per replicate) were conducted in 250-ml glass beakers, under a 16:8 h light:dark cycle in an environmental chamber maintained at $21 \pm 2^\circ\text{C}$. As per the American Society for Testing and Materials method, exposure solutions were not renewed nor were the glochidia fed during the exposure. A composite water sample (10 ml) was collected from each treatment at the beginning of the exposure for determination of dissolved (i.e., filtered through an Acrodisk[™] 0.45 μm in-line-syringe-tip filter; PALL Life Sciences) Cu concentrations in the exposures. Water hardness, pH, dissolved oxygen, major ions (Ca, Mg, Na, K, SO_4 , Cl), and alkalinity were also measured in water samples collected at the beginning each toxicity test. All glassware was washed with 10% nitric acid (Reagent Grade, Fisher Scientific) prior to use and all solutions were made with Millipore[™] water.

Effect of added DOC on copper toxicity

Exposures with L. siliquoidea glochidia. The effect of DOC on the sensitivity of glochidia to Cu was assessed by conducting a series of exposures using reconstituted water and additions of concentrated natural organic matter. Dissolved natural organic matter was collected from Luther Marsh, near Guelph (ON, Canada) ($43^\circ 57'\text{N}$, $80^\circ 26'\text{W}$). One hundred liters of marsh water (raw DOC = 78 mg C/L) was concentrated using a stainless-steel portable reverse-osmosis unit (Limnological Research) to produce a DOC concentrate of 747 mg C/L. In order to displace metals from the binding sites on the DOC and remove them from solution, the concentrate was treated with an H^+ -cation-exchange resin (Amberlite, IR-118H) to a pH of 2.3. The concentrate was stored in the dark, at 4°C until it was used in the Cu toxicity tests, at which time the pH was adjusted to 7.0 using sodium hydroxide (Sigma, Canada).

Individual toxicity tests, referred to as exposures, were conducted at each of 0, 0.5, 1.0, 2.5, 3.75, 5.0, and 7.5 mg DOC/L (nominal) using the concentrated Luther Marsh DOC. Six Cu concentrations (treatments) were assessed in each exposure. Test solutions (water containing DOC and Cu) were created by adding the desired amount of DOC concentrate to 3 L of reconstituted soft water [30]. Then, to each of six 0.5-L aliquots of the DOC water the desired amount of Cu stock solution was added to create a range of Cu concentrations (nominal 0 to 200 μg Cu/L). The Cu-spiked test solutions were held in the dark at 4°C for 48 h before initiation of the exposures to allow the Cu-DOC mixture to equilibrate [31]. Acute toxicity tests were conducted as outlined above.

Single exposure with L. fasciola glochidia. To confirm the relationship between DOC concentration and the sensitivity of glochidia to Cu, we examined the effect of DOC on Cu toxicity

in a second, closely related mussel species, *L. fasciola*. Rather than conducting a series of separate exposures as was done with *L. siliquoides*, a single exposure was conducted at a fixed, acutely toxic level of Cu while the concentration of DOC (i.e., treatments) was varied. A Cu exposure of 18 µg/L was chosen based on a 24-h EC50 of 17.6 µg Cu/L (95% confidence interval [CI] 14.2–22.6) reported by Gillis et al. [11] for this species. Test solutions were created by first adding 18 µg Cu/L (nominal) to 4 L of reconstituted soft water. Then, to each of seven 0.5-L aliquots of the Cu solution, the desired amount of DOC (nominal 0–7.5 mg C/L) was added to create the DOC treatments. The Cu-spiked, DOC exposure solutions were held in the dark at 4°C for 48 h before initiation of the exposures. Acute toxicity tests (including water analysis) were conducted as outlined above. Although the 0 mg/L DOC exposure was the true control in this exposure it contained an acutely toxic concentration of Cu. Therefore, in order to assess the health of *L. fasciola* glochidia (i.e., true control survival) an additional treatment, consisting of only soft water (no Cu or added DOC), was conducted alongside the exposure.

Effect of inherent DOC on copper toxicity

Water was collected from eight locations in southern Ontario, Canada (see Table 1 for locations) using acid-washed 18-L high-density polyethylene buckets. The sites were chosen to represent a gradient of inherent dissolved organic carbon (Table 1). No attempt was made to match other water characteristics (hardness, pH, etc.) between the field sites. In streams and rivers, the water sample was collected from just below the surface in an area where the water was visibly flowing. In lakes, the water was collected from an open water area at least 10 m offshore. Water samples were transported to the laboratory and filtered through a 0.45 µm filter (PALL Life Sciences) and then held in the dark at 4°C until used in exposures (maximum of two weeks). Copper was added (spiked) to each of the field-collected waters to create a range of Cu concentrations (nominal, 0 to 200 µg Cu/L). The Cu-spiked, field-collected waters were held in the dark at 4°C for 48 h before initiation of the exposures. Toxicity tests were conducted in each of the eight Cu-spiked waters with *L. siliquoides* glochidia as outlined above.

Copper analysis

The concentration of dissolved Cu in the test solutions were determined in a filtered (Acrodisc 0.45 µm in-line-syringe-tip filter) water sample collected at initiation ($t = 0$) of the toxicity test. Copper was analyzed using a graphite furnace atomic absorption spectrophotometer (Varian 220FS SpectraAA, Varian Techtron). The detection limit for Cu was 0.02 µg/L. Method blanks (3) and Fisher Scientific calibration standards (every 20 samples) were included in every run. A maximum of 5% difference between duplicates was accepted.

DOC analysis

The concentration of DOC was determined in water samples (10 ml, filtered using an Acrodisc 0.45 µm in-line-syringe-tip filter) collected at the beginning of each exposure using a Shimadzu total organic carbon analyzer (model 5050A; Mandel Scientific). The detection limit for DOC was 0.1 mg C/L. The concentration of total organic carbon in each sample was calculated automatically by subtracting inorganic carbon from total carbon [24].

Statistical analysis

Percent glochidia survival is reported using means with standard errors (SE). The EC50s were determined by Probit Analysis (SPSS v. 11.0) using measured concentrations of dissolved copper ($t = 0$). The EC50s are presented with 95% CIs as EC50 (95% CI). The EC50s were considered to be significantly different when their 95% CI did not overlap [32]. If the 95% CI of two EC50s overlapped, then the Litchfield and Wilcoxon [33] method was applied to determine if they were significantly different. Survival curves were produced using the statistical analysis component (Regression Wizard) of Sigma Plot v. 10.0. Statistical differences ($p < 0.05$) in glochidia survival between treatments in the *L. fasciola* exposure were determined using analysis of variance (ANOVA) followed by Tukey's test.

Comparison of observed EC50 values to BLM predicted water quality criteria

To determine if the current water quality regulations for Cu in North America will protect larval freshwater mussels, the

Table 1. The measured concentration (mg C/L) of dissolved organic carbon (DOC), and observed 24-h copper median effective concentrations (EC50) (95% confidence intervals) in exposures used to assess the effect of DOC on copper toxicity in *L. siliquoides* glochidia; including a series of exposures employing reconstituted soft water augmented with natural DOC (Luther Marsh, ON) and a series of eight natural field-collected waters (no added DOC) from Ontario, Canada. Field sites (water body, site name and location presented) were selected to cover a range of inherent DOC concentrations (5 to 15 mg C/L). The predictions of the U.S. Environmental Protection Agency's copper biotic ligand model (BLM) version (2.2.3) [35] including the site specific final acute values (FAV) and criterion maximum concentrations (CMC) are given for each exposure

Water body	Site name	Latitude	Longitude	DOC (mg C/L)	Observed Cu EC50 (µg/L)	BLM predicted site specific FAV (µg/L)	BLM predicted site specific CMC (µg/L)
Lake Huron	Eagle Harbour	44.18278	-81.62917	5.0	30.9 (26.5-36.3)	100.8	50.4
Lake Saint Clair	Pocket Bay	42.53064	-82.61963	5.7	26.7 (20.3-36.6)	85.7	42.9
Cold Water River	Cold Water	44.71250	-79.64611	6.4	32.1 (21.1-57.1)	135.4	67.7
Saugeen River	Hanover	44.16056	-81.03194	6.9	71.4 (58.3-86.5)	208.6	104.3
Thames River	London	42.97965	-81.12756	9.4	90.9 (77.6-106.7)	211.5	105.8
Grand River	Doon Heritage (Kitchener)	43.40472	-80.4333	11.3	104.4 (97.0-111.0)	315.6	157.8
Sydenham River	Dawn Mills (Dresden)	42.58925	-82.12872	14.7	110.9 (99.7-124.6)	418.1	209.1
Thames River	Thamesford	43.05601	-80.99274	14.8	97.6 (85.9-99.1)	455.9	227.9
Soft water (SW)	—	—	—	0.7	36.1 (30.3-44.0)	5.4	2.7
SW + 0.5 DOC	—	—	—	1.2	42.1 (34.0-53.3)	6.9	3.4
SW + 1.0 DOC	—	—	—	2.0	63.7 (54.2-78.6)	12.5	6.3
SW + 2.5 DOC	—	—	—	3.3	106.5 (85.4-143.3)	32.7	16.4
SW + 3.75 DOC	—	—	—	4.4	150.2 (127.3-187.0)	33.8	16.9

acute toxicity data produced in the present study were compared to Canadian Water Quality Guidelines (CWQG) and U.S. Ambient Water Quality Criteria (U.S. AWQC). Current CWQG are based on hardness of the receiving or testing water [34], whereas U.S. AWQC for Cu are derived using the U.S. EPA Cu biotic ligand model (BLM) [35] and a number of input parameters including temperature, pH, DOC, and major ion concentrations. The online version (2.2.3) of the BLM was used to derive exposure-specific water quality criteria. The measured values for temperature, pH, DOC, Ca, Mg, Na, K, SO₄, Cl, and alkalinity from the exposures (see Table 2) were entered into the BLM program. The BLM output includes a final acute value (FAV) and a criterion maximum concentration (CMC) for each exposure; the BLM-derived values were then compared to the observed 24-h Cu EC50 values determined in this study.

Summary of water quality and mussel distribution data

Raw water quality data collected by the Provincial Water Quality Monitoring Network (PWQMN) were provided by the Ontario Ministry of the Environment (<http://www.ene.gov.on.ca/en/publications/dataproducts>). Copper (μg/L) and DOC (mg C/L) data were provided for water samples taken between 1998 and 2008 covering a total of 134 sites across 10 Ontario conservation authorities (CAs). These 10 CAs, in particular the streams and rivers that flow through them, were selected after consultation with the chair of the Ontario Freshwater Mussel Recovery Team, (T.J. Morris, Canadian Department of Fisheries and Oceans, unpublished data) because they were considered to be significant mussel habitats. The conservation authorities selected for this summary along with their main mussel habitat (i.e., river) included: Ausable Bayfield CA (Ausable River), Grand River CA (Grand River), St. Clair Region CA (Sydenham River), Long Point Region CA (Lyn River), Maitland Valley CA (Maitland River), Quinte CA (Moir and Salmon Rivers), Saugeen Valley CA (Saugeen River), Niagara Peninsula CA (Welland River), and the Upper Thames River CA and Lower Thames Valley CA (Thames River) (Fig. 1). For the purposes of this summary, the water quality and mussel distribution data from the Upper and Lower Thames CAs were combined.

Individual data points (i.e., measured concentrations for either Cu or DOC) were averaged over time for each PWQMN site. For each site, an average 67 individual data points (i.e., chemical analysis conducted on a specific date) were used to

calculate the “site average” for Cu and on average 17 individual data points were used to calculate the site average for DOC. Note that PWQMN did not initiate routine DOC analysis until 2001 or later and that DOC measurements were not taken at all locations (thus the lower number of DOC readings per site compared to Cu). These site averages were then averaged to determine an overall mean for each CA referred herein to as a CA mean. Site averages, rather than individual readings, were used to calculate each CA mean to prevent skewing of the CA mean due to differences in sampling frequency or extreme readings (e.g., unusually high concentrations). The CA range illustrates the lowest and highest individual measured values across the entire CA over the 10 years surveyed. Because the PWQMN copper detection limit was 0.8 μg/L, any readings below this level were set to 0 μg/L for purposes of calculating CA means.

Mussel distribution data were obtained from various sources ranging from individual watershed recovery plans to peer-reviewed literature (Table 3). The total number of living mussel species found in each CA is, to the best of our knowledge, the most up-to-date assessment of mussel distribution in that CA using post-1996 [36] live animal survey data rather than the total number of species that have ever been found in that watershed (i.e., historic data). The number of endangered mussel species reported for each CA was obtained from a series of Department of Fisheries and Oceans (DFO) maps, which present the distribution of aquatic species at risk (fish and mussels) in each CA in Ontario (<http://conservation-ontario.on.ca/projects/DFO.html>).

RESULTS

Effect of added DOC on copper toxicity

L. siliquioidea exposures. The measured concentrations of DOC in the series of *L. siliquioidea* exposures were 0.7, 1.2, 2.0, 3.3, 4.4, 5.2, and 8.7 mg C/L. Control survival (24 h) was >80% in all exposures except in the two highest exposures (5.2 and 8.7 mg C/L), where it was slightly reduced (76 and 78%, respectively). Additions of natural organic matter (Luther Marsh) DOC significantly reduced the toxicity of Cu to *L. siliquioidea* glochidia (Fig. 2). In the control exposure without any added DOC (0.7 mg C/L), the 24-h EC50 was 36.1 (30.3–44.0) μg Cu/L (Table 1). In comparison, the 4.4 mg C/L exposure produced an EC50 of 150.2 (127.3–187.0) μg Cu/L and no EC50 could be determined at the two highest DOC

Table 2. Summary of measured water chemistry including pH, dissolved organic carbon (DOC) (mg/L), and concentrations (mol/L) of calcium, magnesium, sodium, potassium, sulfate, chloride, and dissolved inorganic carbon (DIC) used to predict exposure-specific water quality criteria using the online version (2.2.3) of the U.S. Environmental Protection Agency's copper biotic ligand model (BLM) [35]^a

Site label	pH	DOC (mg/L)	Ca (mol/L)	Mg (mol/L)	Na (mol/L)	K (mol/L)	SO ₄ (mol/L)	Cl (mol/L)	DIC (mol/L)
Soft water	7.45	0.7	2.01E-04	3.04E-04	6.96E-04	3.58E-05	5.22E-04	3.22E-05	6.89E-04
Soft water + 0.5 DOC	7.23	1.2	2.12E-04	3.18E-04	7.48E-04	3.73E-05	5.47E-04	3.44E-05	7.69E-04
Soft water + 1.0 DOC	7.29	2.0	2.25E-04	3.21E-04	7.83E-04	2.92E-05	5.42E-04	3.27E-05	7.57E-04
Soft water + 2.5 DOC	7.62	3.3	2.29E-04	3.60E-04	8.92E-04	4.09E-05	6.30E-04	3.86E-05	8.41E-04
Soft water + 3.75 DOC	7.43	4.4	2.04E-04	3.13E-04	8.05E-04	3.48E-05	5.43E-04	4.01E-05	7.78E-04
Lake Huron	8.30	5.0	6.51E-04	3.18E-04	1.80E-04	2.43E-05	1.71E-04	1.89E-04	1.68E-03
Lake Saint Clair	8.01	5.7	6.59E-04	3.36E-04	2.08E-04	2.56E-05	1.76E-04	2.20E-04	1.42E-03
Cold Water River	8.19	6.4	1.43E-03	6.50E-04	5.05E-04	3.89E-05	1.34E-04	5.27E-04	3.54E-03
Saugeen River	8.54	6.9	1.28E-03	1.13E-03	3.43E-04	2.92E-05	1.72E-04	3.95E-04	4.06E-03
Thames River (London)	8.16	9.4	1.82E-03	8.52E-04	1.58E-03	1.21E-04	6.48E-04	1.76E-03	3.55E-03
Grand River	8.42	11.3	1.26E-03	8.06E-04	1.45E-03	7.67E-05	3.49E-04	1.61E-03	3.11E-03
Sydenham River	8.47	14.7	2.02E-03	5.68E-04	6.52E-04	1.33E-04	4.66E-04	9.17E-04	4.31E-03
Thames River (Thamesford)	8.54	14.8	1.80E-03	9.63E-04	6.87E-04	1.03E-04	5.03E-04	8.49E-04	4.83E-03

^a Temperature was 22°C in all exposures and the concentration of humic acid was set to 10%. Field site locations are given in Table 1.




Fig. 1. Map of southern Ontario, Canada showing location of conservation authorities containing significant freshwater mussel habitats. Copper and dissolved organic carbon concentrations from these areas are summarized in Table 3.

concentrations examined because there was no significant mortality at any of the Cu concentrations tested. The cause of the high mortality observed in 40 μg Cu/L treatment of the 5.2 mg C/L DOC exposure (Fig. 2F) is unknown but an error in exposure solution preparation is suspected. A significant linear relationship ($r^2 = 0.98$, $p = 0.0008$) existed between the concentration of DOC and the resulting Cu EC50 (Fig. 3).

L. fasciola exposure. The 24-h survival of *L. fasciola* glochidia in the soft water treatment without any added DOC or Cu was 97% (± 0.7). The mean measured concentration of Cu in the exposures was 14.9 (± 1.3) $\mu\text{g}/\text{L}$. The measured concentrations of DOC in the treatments were 1.5, 1.9, 2.4, 3.7, 4.7, 5.8, and 7.9 mg C/L. For the control copper treatment, i.e., the treatment that contained an acutely toxic amount of Cu but no added DOC, survival was 44.2% (± 7.7) after 24 h of exposure. A

Table 3. Summary of copper (as total copper) and dissolved organic carbon (DOC) concentrations in significant (stream and river) mussel habitats in southern Ontario and the number of mussel species found in each habitat

Conservation authority	CA mean DOC (mg C/L)	CA range DOC ^a (mg C/L)	CA mean copper ^b ($\mu\text{g}/\text{L}$)	CA range copper ($\mu\text{g}/\text{L}$)	Total no. mussel species [reference]	No. endangered mussel species ^c
Ausable Bayfield	5.8 (1.98), $n = 3$	2.3–13.7	2.02 (0.67), $n = 7$	0–9.6	23 ^d	6 ^c
Grand River	6.0 (0.43), $n = 13$	3.3–13.3	1.64 (0.64), $n = 32$	0–20.1	25 [40]	9
St. Clair Region	6.7 (2.78), $n = 3$	1.9–11.4	2.42 (1.22), $n = 9$	0–12.8	34 [41]	11
Long Point Region	3.6 (NA) ^f , $n = 1$	2.0–10.4	1.27 (NA), $n = 1$	0–3.2	10 ^g	5
Maitland Valley	5.6 (0.56), $n = 5$	3.3–12.1	1.52 (0.42), $n = 13$	0–33.6	9 ^h	2
Quinte	NA	NA	0.97 (0.46), $n = 17$	0–35.7	10 [36]	2
Saugeen Valley	4.7 (1.31), $n = 3$	1.7–9.2	0.59 (0.26), $n = 14$	0–4.3	8 [42]	2
Upper Thames River & Lower Thames Valley	4.9 (0.75), $n = 9$	2.9–11.7	2.08 (1.18), $n = 38$	0–54.3	26 [43]	11
Niagara Peninsula	NA	NA	2.13 (1.18), $n = 3$	0–8.6	10 ⁱ	8

Water quality data (Cu and DOC) was provided by the Provincial Water Quality Monitoring Network (PWQMN) of the Ontario Ministry of the Environment ([39]; <http://www.ene.gov.on.ca/en/publications/dataproducts>). Mean water quality values and ranges (minimum–maximum) are given for data collected from 1998 to 2008. Values reported as mean are the average of all site averages (repeated sampling at one site over time) for each conservation authority (CA). The number of individual site averages used to determine each CA mean or CA range is reported as n , standard deviation is given in parentheses. Mussel distribution data are from various sources (see below).

^a PWQMN did not initiate routine DOC analysis until 2001 or later, and DOC measurements were not taken at all locations.

^b The PWQMN copper detection limit was 0.8 $\mu\text{g}/\text{L}$, therefore any copper concentration below this level has been reported as 0 $\mu\text{g}/\text{L}$.

^c Designated as such by the Committee on the Status of Endangered Wildlife in Canada.

^d Ausable River Recovery Team. 2005. Recovery strategy for species at risk in the Ausable River (<http://www.abca.on.ca/publications.php>).

^e Endangered species data are from the Canadian Department of Fisheries and Oceans (<http://conservation-ontario.on.ca/projects/DFO.html>).

^f NA = data not available.

^g J.L. Metcalfe-Smith, S.K. Staton, Environment Canada, Burlington, ON, Canada, unpublished data.

^h D.J. McGoldrick, J.L. Metcalfe-Smith, Environment Canada, Burlington, ON, Canada, unpublished data.

ⁱ Todd Morris, Department of Fisheries and Oceans, Burlington, ON, Canada, unpublished data.




Fig. 2. Percent survival of *Lampsilis siliquoidea* glochidia when exposed (24 h) to copper in reconstituted soft water with varying amounts of added dissolved organic carbon (DOC). (A) 0.7 mg C/L, (B) 1.2 mg C/L, (C) 2.0 mg C/L, (D) 3.3 mg C/L, (E) 4.4 mg C/L, and (F) 5.2 mg C/L. Error bars represent standard errors around a mean of three replicates. EC50 = median effective concentration.

significant increase in glochidia survival (compared to the control) was observed when concentrated Luther Marsh DOC was added to the Cu exposures. Glochidia survival in all DOC augmented treatments was more than 80%, with no significant difference observed between the treatments (1.9–7.9 mg C/L).

Effect of inherent DOC on copper toxicity

The concentration of naturally occurring (i.e., inherent) DOC in the field-collected waters are given in Table 1. Control

survival ranged from 86 to 95% in the eight field-collected waters. The concentration of DOC had a significant effect on the sensitivity of *L. siliquoidea* glochidia to waterborne Cu exposure (Fig. 4). The 24-h EC50 values ranged from 30.9 µg Cu/L in water collected from Lake Huron (DOC, 5.0 mg C/L) to 110.9 µg Cu/L in water collected from the Sydenham River (DOC, 14.7 mg C/L). Regression analysis revealed a significant relationship ($r^2 = 0.79$, $p = 0.0031$) between the concentration of inherent DOC and the resulting Cu EC50 (Fig. 4).




Fig. 3. The relationship between the concentration of dissolved organic carbon (DOC) and the resulting copper median effective concentration (EC50) of *Lampsilis siliquoidea* glochidia. Each point represents a copper 24-h median effective concentration (EC50) with 95% confidence intervals as error bars. Each EC50 was derived from a toxicity test comprised of six copper treatments, each with three replicates. EC50s indicated with an asterisk are significantly elevated above the control EC50 (no DOC added).

DISCUSSION

Effect of added DOC on copper toxicity

The significant linear relationship between the concentration of DOC (as Luther Marsh DOC) in the exposure water and the resulting Cu EC50 illustrates the significant ameliorating effect that natural DOC has on Cu toxicity in glochidia. An increase in the concentration of DOC from 0.7 mg/L to 4.4 mg/L resulted in a fourfold increase (36–150 µg Cu/L) in the EC50 of *L. siliquoidea* glochidia. Similarly, a significant increase in the survival of *L. fasciola* glochidia was observed when natural DOC was added to an acutely toxic Cu exposure. Negatively charged ligands such as DOC bind with positively charged cations such as Cu, reducing the amount of free Cu in solution and thus reduce the bioavailability of Cu [16]. The ameliorating effect of natural DOC on Cu toxicity in juvenile freshwater mussels (*L. siliquoidea*) was recently demonstrated by Wang et al. [26]. They also reported a significant linear relationship




Fig. 4. The relationship between the concentration of inherent (i.e., not added) dissolved organic carbon (DOC) in various copper-spiked, field-collected waters (see Table 1 for site details and Table 3 for water composition) and the resulting median effective concentration (EC50) (95% confidence intervals as error bars) for *Lampsilis siliquoidea* glochidia. Each point represents an EC50 derived from a toxicity test (six copper concentrations, each with three replicates) conducted in water collected from one of eight different field locations.

between the concentration of DOC in an exposure and the Cu EC50. In one series of exposures with additions of Luther Marsh DOC and reconstituted hard water, a 4.2-fold increase in the juvenile mussel Cu EC50 was observed when DOC was raised from 0 to 5 mg C/L [26]. This is nearly identical to the fourfold increase in the glochidia Cu EC50 reported in the present study when DOC was increased from <1 to 4.4 mg C/L using Luther Marsh DOC and reconstituted soft water. Additions of DOC have also been shown to reduce Cu toxicity in a number of aquatic species. In cladocerans, Giesy et al. [20] reported that the presence of organic matter reduced the accumulation of Cu in *Simocephalus serrulatus*, and De Schampelaere and Janssen [21], as well as De Schampelaere et al. [22] demonstrated the protective effect of natural DOC additions on the copper toxicity in *Daphnia magna*. Similarly, in fish (rainbow trout [24]; fathead minnow [18]) acute Cu toxicity has been shown to decrease when the concentration of DOC in the exposure water was increased. Nadella et al. [37] also recently demonstrated the ameliorating effect of added DOC on the Cu sensitivity of the embryos of marine mussels (*Mytilus trossolus*).

Effect of inherent DOC on copper toxicity

The concentration of inherent DOC in natural waters had a significant effect on the sensitivity of glochidia to Cu. Copper sensitivity of *L. siliquoidea* glochidia exposed to Cu-spiked water from Lake Huron (EC50 30.9 µg/L), the field-collected water with the lowest concentration of DOC (5.0 mg C/L), was similar to the EC50 (36.1 µg Cu/L) in reconstituted soft water exposures; however, when acute copper toxicity was assessed in water collected from the Grand River (DOC 11.3 mg C/L), the resulting EC50 was nearly three times higher than that of the soft water exposure (EC50 104.4 µg Cu/L). A significant relationship between the concentration of inherent DOC and the toxicity of Cu was also demonstrated in the embryos of marine mussels [25]. Arnold [25] found that the Cu EC50 of *Mytilus galloprovincialis* was highly correlated ($r^2 = 0.71, p < 0.001$) to the concentration of inherent DOC in the water (54 natural marine waters tested).

It should be noted that dissolved organic carbon was not the only water chemistry parameter that varied across the natural waters employed in the present study; water hardness also ranged from 84 to 284 mg CaCO₃/L (Table 2 presents water composition of the field-collected waters). Interestingly, there was a positive and significant correlation (Pearson correlation coefficient 0.81, $p = 0.015$) between water hardness and the concentration of inherent DOC in the natural waters used in the present study. The bioavailability of Cu is known to be influenced by a number of water chemistry parameters, including the concentrations of major cations (Ca, Mg, Na) [14,15]. Water hardness has been shown to decrease Cu sensitivity of glochidia in exposures using reconstituted waters. For example *Epioblasma triquetra* glochidia exposed to Cu in moderately hard (166 mg CaCO₃/L) water had an EC50 twofold higher than those exposed in soft (40 mg CaCO₃/L) water [11]. Although the effect of water hardness is significant, increases in water hardness do not appear to affect Cu bioavailability as strongly as increases in DOC. Again, our earlier exposures with *E. triquetra* and Aldrich humic acid demonstrated that a small increase in the DOC also resulted in a doubling of the EC50 [11]. However, the presence of other toxicity modifying factors in our field-collected water exposures explains the variability observed in the linear regression of DOC on EC50 for natural waters (Fig. 4) versus reconstituted waters (Fig. 2).

Comparison of added DOC and inherent DOC

On a per concentration basis, it appears that the Luther Marsh DOC, which was added to toxicity tests using reconstituted water, was much more effective at reducing the bioavailability of Cu than the natural DOC in the field-collected waters. For example, exposures with 4.4 mg C/L of Luther Marsh DOC resulted in an EC50 of 150 $\mu\text{g Cu/L}$. In comparison, the EC50 values in Cu-spiked natural waters with inherent DOC concentrations of 5 to 6 mg C/L all had EC50 values less than 40 $\mu\text{g Cu/L}$. A number of factors may have contributed to this difference including the source of the DOC. Schwartz et al. [24] compared natural organic matter (NOM) from a number of sources with respect to their ameliorating effect on metal toxicity. They demonstrated that the NOM from Luther Marsh, an optically dark NOM which is dominated by allochthonous (terrestrially derived) material, reduced Ag toxicity in fish to a greater degree than did other optically lighter NOMs. Typically, darker colored NOMs have more metal complexing abilities than the lighter autochthonous NOMs (i.e., those derived from aquatic photosynthesis) that are found in large lakes and oceans [23,24]. In fact, Luider et al. [23] reported that darker colored NOMs had about twofold greater Cu-binding capacity compared to the lighter colored ones.

Although we did not isolate the natural organic matter from the field-collected waters to quantify darkness (or composition) in the present study, based on earlier research it seems appropriate that the DOC from the large lakes and rivers we tested would be more autochthonous in nature (therefore less effective at binding Cu) and this may explain the relatively lower protective effects compared to Luther Marsh DOC. An alternative, or perhaps confounding, reason for the observed difference between the ameliorating capabilities of the Luther Marsh and other field-collected DOCs, may lie in the differences in the composition of the water used in the exposures. The reconstituted water used in the Luther Marsh exposures was much softer (40 mg CaCO_3/L) than the natural waters used to assess the effect of inherent DOC (84–284 mg CaCO_3/L). As discussed above, glochidia have been shown to be less sensitive to Cu when exposed in hard water compared to soft water [9,11]. The increased concentration of major ions (Ca^{2+} , Mg^{+}) in hard water leads to increased competition between those ions and the copper ion (Cu^{2+}) for binding sites on the organism [14,15]. In addition to competition for binding sites on the organism, there will also be competition among the various cations (Ca^{2+} , Mg^{+} , Cu^{2+}) for binding sites on the DOC molecules. Mantoura et al. [19] reported that Mg and Ca compete with Cu for binding sites on the humic acid molecule. Similarly, Iglesias et al. [38] found that the degree of Cu complexation with fulvic acid decreased as the concentration of Ca^{2+} in the exposure medium increased. This would indicate that DOC would be more effective at reducing the bioavailability of Cu in soft water than in hard water. Therefore, we suggest that the enhanced protective ability of the Luther Marsh DOC in soft waters compared to the DOC from field collected waters is a result of both the increased metal-binding capabilities of this allochthonous dominated DOC and the ion-dilute soft water used in the exposures.

Copper sensitivity: water quality criteria and guidelines

Currently, Canadian Water Quality Guidelines for Cu are based on water hardness. For waters with hardness less than 120 mg CaCO_3/L the guideline is 2 $\mu\text{g Cu/L}$ [34]. For waters with hardness between 120 and 180 mg CaCO_3/L , the guideline

is 3 $\mu\text{g Cu/L}$ [34]. These guidelines would be protective considering the EC50 values derived from the natural water exposures of the present study ranged from 27 to 111 $\mu\text{g Cu/L}$. Currently, the most scientifically advanced regulatory approach for Cu is the U.S. EPA Ambient Water Quality Criteria, which are derived on a site-specific basis using the biotic ligand model (BLM) [35]. Table 1 presents the observed EC50 values alongside the BLM predicted final acute values (FAVs) and criterion maximum concentrations (CMC) derived for the exposures in the present study. The CMC (or acute criteria) values (half the FAV) produced range between 43 and 228 $\mu\text{g Cu/L}$ (Table 2). Typically, the actual water quality criteria, the criterion continuous concentration (CCC or chronic criteria), which is the maximum 4-d mean concentration that is allowed over a three-year period, is derived by applying the acute-to-chronic ratio to the CMC. The application of the BLM (v. 2.2.3, in Cu WQC Calculation mode) produced predicted FAV values that for every natural field-collected water were greater than the measured toxicity (Table 1). A number of explanations are possible. For example, it is possible that the natural waters came with a contaminant load that resulted in glochidia being sensitized to Cu. Alternatively, it is possible that DOC bound Cu is bioavailable to glochidia and contributes to toxicity. It is also possible that BLM predictions generally overestimate protective effects, particularly at elevated pH. All of the natural waters tested had pH values over 8.0 (range 8.1–8.6). In any event, the BLM was not able to predicted Cu toxicity to glochidia in natural waters and set FAV values that were inappropriate. Whether the anomaly between measured and BLM predicted toxicity results from unique features of the organism or the model (or both) is unknown; however, it is deserving of further study.

Implications for native populations of mussels

The rivers and streams of the lower Great Lakes Basin contain the richest assemblage of freshwater mussel species in Canada [36]. Forty species of mussels are found in this area, including 11 federally endangered species (<http://conservation-ontario.on.ca/projects/DFO.html>). After surveying both historic (pre-1960) and more recent mussel distribution data of southern Ontario, Metcalfe-Smith et al. [36] concluded that significant species losses and community changes in the freshwater mussel populations have occurred. They suggested that even by conservative estimates, somewhere between one-sixth and one-third of the mussel fauna in the Grand, Thames, and Sydenham Rivers have been lost. Although the decline of native mussels in North America has been attributed to a number of factors, including habitat alternation, loss of fish hosts, and the invasive zebra mussel, environmental pollution is also considered to have contributed to the decline of native freshwater mussels. However, until recently research in this area has been somewhat limited and thus the impact of waterborne contaminants on native mussel populations is still uncertain. In fact, all of the recovery strategies for freshwater mussel produced to date in Canada have indicated the "urgent need" to examine the threat that waterborne contaminants pose to the recovery of imperiled freshwater mussels. Therefore, to address this need, we examined the results of this laboratory-based toxicity study in the context of the measured levels of Cu in the mussel's natural habitat.

A summary of the copper (as total Cu) and DOC concentrations found in significant (stream and river) mussel habitats in Ontario over the past 10 years is presented in Table 3. Data were compiled by conservation authority. The number of

mussel species found within each conservation authority area, as well as the number of endangered species, are also included. According to PWQMN data, the concentration of Cu found in mussel habitats in southern Ontario has ranged from less than detectable ($<0.8 \mu\text{g/L}$) to $54.3 \mu\text{g/L}$ over the past 10 years, and the mean Cu concentration (see *Materials and Methods* for explanation) across the CAs ranged from 0.6 to $2.4 \mu\text{g Cu/L}$ (Table 3). Although the Ontario Ministry of the Environment did not initiate routine DOC analysis until 2001, and DOC measurements were not taken at all locations, the data available indicate that the concentration of inherent DOC ranged from 1.9 to 13.7 mg C/L at these sites. The overall CA means for DOC spanned 3.6 to 6.7 mg C/L (Table 3). Therefore, according to our earlier [11] laboratory-based acute toxicity tests using reconstituted soft water (no added DOC), the mean Cu concentration in the mussel habitats surveyed does not exceed the level found to be acutely toxic to even the most sensitive of the seven species tested (*V. fabalis* 24 h EC₅₀, $6.9 \mu\text{g Cu/L}$). This conclusion is supported by EC₅₀ values derived from toxicity tests conducted in Cu-spiked natural waters in the present study. For example, waters with similar levels of inherent DOC as the mean levels found in the CAs examined (4 to 7 mg C/L) produced EC₅₀ values that ranged from 31 to $71 \mu\text{g Cu/L}$. This is more than an order of magnitude above the mean level of Cu reported in the rivers examined. Even though the mean levels of DOC and Cu in the mussel habitats of southern Ontario indicate that exposure to acutely toxic levels of Cu is not the typical situation, the episodic pulses or spikes of Cu may be of concern, depending on the DOC concentration of the water at that time. For instance, in the Thames River, a habitat that supports 11 federally endangered species of mussels, Cu has been found to exceed $50 \mu\text{g/L}$ on occasion. With variation in DOC concentrations (for example, 2.9 – 11.7 mg C/L in the Thames River, PWQMN data) even short-term spikes in Cu concentration could be harmful, especially if the spike occurs at a sensitive stage in the reproductive cycle, particularly during the time that glochidia are released into the water column (May–October). The significant mussel habitats of southern Ontario are typically DOC-rich and while there are many instances in nature in which excess nutrients and DOC can lead to a deterioration of water quality (i.e., eutrophication), the results of the present study indicate that the inherent DOC found in mussel habitats in southern Ontario should, on average, provide freshwater mussels with protection from acute Cu toxicity, although further study into the effects of episodic Cu release on endemic freshwater mussels would be beneficial.

Although the acute Cu sensitivity of glochidia has been assessed in a variety of mussel species [8,10,11], the diversity of the exposure water used in those studies was very limited. Typically, reconstituted waters have been used to determine an organism's sensitivity to a contaminant and, although such standardized waters are useful for comparing sensitivities among species and among studies, they do not necessarily reflect the complexity of the water found in an organism's natural habitat. Because freshwater mussels are found in a wide range of habitats from small urbanized streams to large oligotrophic lakes, it is important to understand their vulnerability to contaminants under such diverse conditions. By assessing the Cu sensitivity of glochidia in a range of natural waters we have demonstrated that the sensitivity of glochidia to Cu not only depends on the species [11], but also on the composition of the water, especially the concentration of DOC, in which they live.

CONCLUSION

In conclusion, the present study has demonstrated the significant protective effect that natural organic matter (as DOC) has on acute Cu toxicity in glochidia. The significant relationship between the concentration of inherent DOC and copper toxicity emphasizes that an understanding of the DOC level in the mussel's natural habitat is necessary to assess the bioavailability of Cu and its potential threat to the sensitive early life stages of freshwater mussels.

Acknowledgement—This work was supported by the World Wildlife Fund (Endangered Species Recovery Fund (WWF-Canada)), the Copper Development Association, and the Species at Risk – Espèces en Péri (Department of Fisheries and Oceans Canada). The authors thank R. Mitchell, K. McNichols, A. Schwalb, P. Ragaz, J. Klink, O. Birceanu, and G. Nishihara for assistance in the laboratory and field, as well as A. Todd and S. Sunderani of the Ontario Ministry of the Environment for providing the raw water quality data (PWQMN), and T. Morris (DFO) for assistance selecting mussel habitats and making mussel distribution data available to us. O. Bireanu was supported by an NSERC Metals in the Human Environment Strategic Network grant.

REFERENCES

1. U.S. Fish and Wildlife Service 1990. Endangered and threatened wildlife and plants. *Fed Reg* 59:58982–59028.
2. Williams JD, Warren ML, Cummings KS, Harris JL, Neves RJ. 1993. Conservation status of freshwater mussels of the United States and Canada. *Fisheries* 18:6–22.
3. Bogan AE. 1993. Freshwater bivalve extinctions (Mollusca: Unionidae): a search for causes. *Am Zool* 33:599–609.
4. Gillis PL, Mackie GL. 1994. Impact of the zebra mussel, *Dreissena polymorpha*, on populations of Unionidae (Bivalvia) in Lake St. Clair. *Can J Zool* 72:1260–1271.
5. Fleming WJ, Augspurger TP, Alderman JA. 1995. Freshwater mussel die-off attributed to anticholinesterase poisoning. *Environ Toxicol Chem* 14:877–879.
6. Huebner JD, Pynnönen KS. 1992. Viability of two species of *Anodonta* exposed to low pH and selected metals. *Can J Zool* 70:2348–2355.
7. Cope WG, Bringolf RB, Buchwalter DB, Newton TJ, Ingersoll CG, Wang N, Augspurger T, Dwyer FJ, Barnhart MC, Neves RJ, Hammer E. 2008. Differential exposure, duration, and sensitivity of unionoidean bivalve life stages to environmental contaminants. *J North Am Benthol Soc* 27:451–462.
8. Jacobson PJ, Neves RJ, Cherry DS, Farris JL. 1997. Sensitivity of glochidial stages of freshwater mussels (Bivalvia: Unionidae) to copper. *Environ Toxicol Chem* 16:2384–2392.
9. Keller AE, Zam SG. 1991. The acute toxicity of selected metals to the freshwater mussel, *Anodonta imbellilis*. *Environ Toxicol Chem* 10:539–546.
10. Wang N, Ingersoll CG, Hardesty DK, Ivey CD, Kunz JL, May TW, Dwyer FJ, Roberts AD, Augspurger T, Kane CM, Neves RJ, Barnhart MC. 2007. Acute toxicity of copper, ammonia, and chlorine to glochidia and juvenile mussels. *Environ Toxicol Chem* 26:2036–2047.
11. Gillis PL, Mitchell RJ, Schwalb AN, McNichols KA, Mackie GL, Wood CM, Ackerman JD. 2008. Sensitivity of the glochidia (larvae) of freshwater mussels to copper: Assessing the effect of water hardness and dissolved organic carbon on the sensitivity of endangered species. *Aquat Toxicol* 88:137–145.
12. March FA, Dwyer FJ, Augspurger T, Ingersoll CG, Wang N, Mebane CA. 2007. An evaluation of water quality guidance and standards for copper. *Environ Toxicol Chem* 26:2066–2074.
13. Ward S, Augspurger T, Dwyer FJ, Kane C, Ingersoll CG. 2007. Risk assessment of water quality in three North Carolina streams supporting federally endangered freshwater mussels. *Environ Toxicol Chem* 26:2075–2085.
14. Pagenkopf GK. 1983. Gill surface interaction model for trace-metal toxicity to fishes: Role of complexation, pH, and water hardness. *Environ Sci Technol* 17:342–347.
15. Playle RC. 1998. Modelling metal interactions at fish gills. *Sci Total Environ* 219:147–163.
16. Playle RC, Dixon DG, Burnison K. 1993. Copper and cadmium binding to fish gills: Modification by dissolved organic carbon and synthetic ligands. *Can J Fish Aquat Sci* 50:2667–2677.
17. Campbell PGC. 1995. Interactions between trace metals and aquatic organisms: A critique of the free-ion activity model. In Tessier A, Turner

- DR, eds *Metal Speciation and Bioavailability in Aquatic Systems*. John Wiley, New York, NY, USA, pp 45-102.
18. Santore RC, Di Toro DM, Paquin PR, Allen HE, Meyer JS. 2001. A biotic ligand model of the acute toxicity of metals: II: Application to acute copper toxicity in freshwater fish and *Daphnia*. *Environ Toxicol Chem* 20:2397-2402.
 19. Mantoura RFC, Dickson A, Riley JP. 1978. The complexation of metals with humic materials in natural waters. *Estuar Coast Mar Sci* 6:386-408.
 20. Giesy JP, Newell A, Leversee GL. 1983. Copper speciation in soft, acid, humic waters: effects on copper bioaccumulation by and toxicity to *Simocephalus serrulatus* (Daphnidae). *Sci Total Environ* 28:23-36.
 21. De Schamphelaere KAC, Janssen CR. 2004. Effects of dissolved organic carbon concentration and source, pH, and water hardness on chronic toxicity of copper to *Daphnia magna*. *Environ Toxicol Chem* 23:115-1122.
 22. De Schamphelaere KAC, Vasconcelos FM, Tack FMG, Allen HE, Janssen CR. 2004. The effect of dissolved organic matter source on acute copper toxicity of *Daphnia magna*. *Environ Toxicol Chem* 23:1248-1255.
 23. Luider CD, Crusius J, Playle RC, Curtis PJ. 2004. Influence of natural organic matter source on copper speciation as demonstrated by Cu binding to fish gills, by ion selective electrode, and by DGT gel sampler. *Environ Sci Technol* 38:2865-2872.
 24. Schwartz ML, Curtis PJ, Playle RC. 2004. Influence of natural organic matter source on acute copper, lead, and cadmium toxicity to rainbow trout (*Oncorhynchus mykiss*). *Environ Toxicol Chem* 23:2889-2899.
 25. Arnold WR. 2005. Effects of dissolved organic carbon on copper toxicity: Implications for saltwater copper criteria. *Integr Environ Assess Manag* 1:34-39.
 26. Wang N, Mebane CA, Kunz JL, Ingersoll CG, May TW, Arnold WR, Santore RC, Augspurger T, Dwyer FJ, Barnhart MC. 2009. Evaluation of acute copper toxicity to juvenile freshwater mussels (fatmucket, *Lampsilis siliquoidea*) in natural and reconstituted waters. *Environ Toxicol Chem* 28:2367-2377.
 27. Markich SJ, Brown PL, Jeffree RA, Lim RP. 2003. The effects of dissolved organic carbon on the toxicity of cadmium and copper to a freshwater bivalve: further support for the extended free ion activity model. *Arch Environ Contam Toxicol* 45:479-491.
 28. Hanstén C, Heino M, Pynnönen K. 1996. Viability of glochidia of *Anodonta anatine* (Unionidae) exposed to selected metals and chelating agents. *Aquat Toxicol* 34:1-12.
 29. American Society for Testing and Materials. 2006. Standard guide for conducting laboratory toxicity tests with freshwater mussels. E2455-06. In *Annual Book of ASTM Standards*, Vol 11.06. West Conshohocken, PA, pp 1393-1444.
 30. American Society for Testing and Materials. 2006. Standard guide for conducting acute toxicity tests on test materials with fishes, macro-invertebrates and amphibians. E 729-96. In *Annual Book of ASTM Standards*, Vol 11.06. West Conshohocken, PA, pp 79-100.
 31. Ma H, Kim SD, Cha DK, Allen HE. 1999. Effect of kinetics of complexation by humic acid on toxicity of copper to *Ceriodaphnia dubia*. *Environ Toxicol Chem* 18:828-837.
 32. Environment Canada. 2005. Guidance document on statistical methods for environmental toxicity tests. EPS 1/RM/46. Method Development and Applications Section, Environmental Technology Centre, Environment Canada, Ottawa, ON.
 33. Litchfield JT, Wilcoxon F. 1949. A simplified method of evaluating dose-effective experiments. *J Pharmacol Exp Ther* 96:99-113.
 34. Canadian Council of Ministers of the Environment. 2005. *Canadian water quality guidelines*. Environment Canada, Ottawa, ON.
 35. U.S., Environmental Protection Agency. 2007. Aquatic life ambient freshwater quality criteria-Copper. EPA-822-R-07-001. Office of Water, Washington, DC.
 36. Metcalfe-Smith JL, Staton SK, Mackie GL, Lane NM. 1998. Changes in the biodiversity of freshwater mussels in the Canadian waters of the lower Great Lakes drainage basin over the past 140 years. *J Gt Lakes Res* 24:845-858.
 37. Nadella SR, Fitzpatrick JL, Franklin N, Bucking C, Smith S, Wood CM. 2009. Toxicity of dissolved Cu, Zn, Ni and Cd to developing embryos of the blue mussel (*Mytilus trossolus*) and the protective effect of dissolved organic carbon. *Comp Biochem Physiol C* 149:340-348.
 38. Iglesias A, Lopez R, Fiol S, Antelo JM, Arce F. 2003. Analysis of copper and calcium-fulvic acid complexation and competition effects. *Water Res* 37:3749-3755.
 39. Ontario Ministry of the Environment. 2009. *Provincial Water Quality Monitoring Network (PWQMN) Database*. Toronto, ON, Canada.
 40. Metcalfe-Smith JL, Mackie GL, Di Maio J, Staton SK. 2000. Changes over time in the diversity and distribution of freshwater mussels (Unionidae) in the Grand River, Southwestern Ontario. *J Gt Lakes Res* 26:445-459.
 41. Jacques Whitford Environment Limited. 2001. Sydenham River recovery project: Synthesis and analysis of background data. Project ONO50368. Ottawa, ON, Canada.
 42. Morris TJ, Granados M, Edwards A. 2007. A preliminary survey of the freshwater mussels of the Saugeen River Watershed, Ontario. Manuscript 2809. Canadian Manuscript Report of Fisheries and Aquatic Sciences, Ottawa, ON.
 43. Morris TJ, Edwards A. 2007. Freshwater mussel communities of the Thames River, Ontario: 2004-2005. Manuscript 2810. Canadian Manuscript Report of Fisheries and Aquatic Sciences, Ottawa, ON.